

Comparability in Bruhat Orders

Jonathan Boretsky (McGill University / LACIM)

joint with Alvaro Cornejo, Reuven Hodges, Paul Horn, Nathan Lesnevich, Tyrrell McAllister

Permutation Patterns, Cal State LA · July 23, 2026

LACIM

Laboratoire d'algèbre, de combinatoire et
d'informatique mathématique



A word cloud where the word 'permutation' is the largest and most central. Other words of varying sizes and colors (blue, green, yellow, red) are scattered around it, including 'patterns', 'set', 'function', 'pumber', 'avoiding', 'classes', 'denote', 'numbers', 'words', and 'functions'.



Permutation Orders in Nature

In studying permutations, there are two natural partial orders on $\mathfrak{S}_n$ which emerge in many interesting ways:

Permutation Orders in Nature

In studying permutations, there are two natural partial orders on $\mathfrak{S}_n$ which emerge in many interesting ways:

Sorting & inversions

- A permutation is determined by its inversions. Inclusion of inversions defines a poset on permutations.
- The one-skeleton of the **permutohedron** can be viewed as a Hasse diagram for a poset on permutations. [Ziegler '95].
- Say $\sigma \preceq_W \tau$ if σ appears in the process of applying a **bubble sort** to τ . [Knuth '98]

Permutation Orders in Nature

In studying permutations, there are two natural partial orders on $\mathfrak{S}_n$ which emerge in many interesting ways:

Sorting & inversions

- A permutation is determined by its inversions. Inclusion of inversions defines a poset on permutations.
- The one-skeleton of the **permutohedron** can be viewed as a Hasse diagram for a poset on permutations. [Ziegler '95].
- Say $\sigma \preceq_W \tau$ if σ appears in the process of applying a **bubble sort** to τ . [Knuth '98]

Permutation Orders in Nature

In studying permutations, there are two natural partial orders on $\mathfrak{S}_n$ which emerge in many interesting ways:

Sorting & inversions

- A permutation is determined by its inversions. Inclusion of inversions defines a poset on permutations.
- The one-skeleton of the **permutohedron** can be viewed as a Hasse diagram for a poset on permutations. [Ziegler '95].
- Say $\sigma \preceq_W \tau$ if σ appears in the process of applying a **bubble sort** to τ . [Knuth '98]

Permutation Orders in Nature

In studying permutations, there are two natural partial orders on $\mathfrak{S}_n$ which emerge in many interesting ways:

Sorting & inversions

- A permutation is determined by its inversions. Inclusion of inversions defines a poset on permutations.
- The one-skeleton of the **permutohedron** can be viewed as a Hasse diagram for a poset on permutations. [Ziegler '95].
- Say $\sigma \preceq_W \tau$ if σ appears in the process of applying a **bubble sort** to τ . [Knuth '98]

$\rightsquigarrow$ These all give **weak** Bruhat order.

Permutation Orders in Nature

In studying permutations, there are two natural partial orders on $\mathfrak{S}_n$ which emerge in many interesting ways:

Sorting & inversions

- A permutation is determined by its inversions. Inclusion of inversions defines a poset on permutations.
- The one-skeleton of the **permutohedron** can be viewed as a Hasse diagram for a poset on permutations. [Ziegler '95].
- Say $\sigma \preceq_W \tau$ if σ appears in the process of applying a **bubble sort** to τ . [Knuth '98]

$\rightsquigarrow$ These all give **weak** Bruhat order.

Geometry & subwords

- Schubert cells Ω_σ° in GL_n/B are indexed by permutations. Say $\sigma \preceq_B \tau$ if the closure of Ω_τ° contains Ω_σ° . [Ehresmann '34; Chevalley '94]
- Say $\sigma \preceq_B \tau$ if a reduced word for σ is a *subword* of a reduced word for τ . [Björner–Brenti '05]
- Dominance of permutation matrices defines a poset on permutations.

Permutation Orders in Nature

In studying permutations, there are two natural partial orders on $\mathfrak{S}_n$ which emerge in many interesting ways:

Sorting & inversions

- A permutation is determined by its inversions. Inclusion of inversions defines a poset on permutations.
- The one-skeleton of the **permutohedron** can be viewed as a Hasse diagram for a poset on permutations. [Ziegler '95].
- Say $\sigma \preceq_W \tau$ if σ appears in the process of applying a **bubble sort** to τ . [Knuth '98]

$\rightsquigarrow$ These all give **weak** Bruhat order.

Geometry & subwords

- Schubert cells Ω_σ° in GL_n/B are indexed by permutations. Say $\sigma \preceq_B \tau$ if the closure of Ω_τ° contains Ω_σ° . [Ehresmann '34; Chevalley '94]
- Say $\sigma \preceq_B \tau$ if a reduced word for σ is a *subword* of a reduced word for τ . [Björner–Brenti '05]
- Dominance of permutation matrices defines a poset on permutations.

Permutation Orders in Nature

In studying permutations, there are two natural partial orders on $\mathfrak{S}_n$ which emerge in many interesting ways:

Sorting & inversions

- A permutation is determined by its inversions. Inclusion of inversions defines a poset on permutations.
- The one-skeleton of the **permutohedron** can be viewed as a Hasse diagram for a poset on permutations. [Ziegler '95].
- Say $\sigma \preceq_W \tau$ if σ appears in the process of applying a **bubble sort** to τ . [Knuth '98]

$\rightsquigarrow$ These all give **weak** Bruhat order.

Geometry & subwords

- Schubert cells Ω_σ° in GL_n/B are indexed by permutations. Say $\sigma \preceq_B \tau$ if the closure of Ω_τ° contains Ω_σ° . [Ehresmann '34; Chevalley '94]
- Say $\sigma \preceq_B \tau$ if a reduced word for σ is a *subword* of a reduced word for τ . [Björner–Brenti '05]
- Dominance of permutation matrices defines a poset on permutations.

Permutation Orders in Nature

In studying permutations, there are two natural partial orders on $\mathfrak{S}_n$ which emerge in many interesting ways:

Sorting & inversions

- A permutation is determined by its inversions. Inclusion of inversions defines a poset on permutations.
- The one-skeleton of the **permutohedron** can be viewed as a Hasse diagram for a poset on permutations. [Ziegler '95].
- Say $\sigma \preceq_W \tau$ if σ appears in the process of applying a **bubble sort** to τ . [Knuth '98]

$\rightsquigarrow$ These all give **weak** Bruhat order.

Geometry & subwords

- Schubert cells Ω_σ° in GL_n/B are indexed by permutations. Say $\sigma \preceq_B \tau$ if the closure of Ω_τ° contains Ω_σ° . [Ehresmann '34; Chevalley '94]
- Say $\sigma \preceq_B \tau$ if a reduced word for σ is a *subword* of a reduced word for τ . [Björner–Brenti '05]
- Dominance of permutation matrices defines a poset on permutations.


$\rightsquigarrow$ These all give **strong** Bruhat order.

Weak Bruhat order

Write $\tau = s_{i_1} \cdots s_{i_k}$ as a *reduced* product of simple transpositions.

Definition (weak order)

$\sigma \preceq_W \tau$ iff σ is a **prefix** of some reduced word for τ : $\sigma = s_{i_1} \cdots s_{i_m}$, $m \leq k$.




Weak Bruhat order

Write $\tau = s_{i_1} \cdots s_{i_k}$ as a *reduced* product of simple transpositions.

Definition (weak order)

$\sigma \preceq_W \tau$ iff σ is a **prefix** of some reduced word for τ : $\sigma = s_{i_1} \cdots s_{i_m}$, $m \leq k$.

- Let $\text{Inv}(\sigma)$ be the set of pairs (i, j) such that $i < j$ and $\sigma(i) > \sigma(j)$. Then, [Björner–Wachs '91]:
 $\sigma \preceq_W \tau \iff \text{Inv}(\sigma) \subseteq \text{Inv}(\tau)$.
- Going *up* an edge = swapping an adjacent pair into an *inversion*.
- Cover relations: an occurrence of the mesh pattern  becomes an occurrence of the mesh pattern .



$\{ij\}$ next to σ is $\text{Inv}(\sigma)$

A basic question about any finite poset

Given a poset $(P, \preceq)$, pick two elements independently and uniformly at random.

How likely are they comparable? $\mathbb{P}(X \preceq Y) = \frac{\#\{\text{intervals}\}}{|P|^2}.$

A basic question about any finite poset

Given a poset $(P, \preceq)$, pick two elements independently and uniformly at random.

How likely are they comparable? $\mathbb{P}(X \preceq Y) = \frac{\#\{\text{intervals}\}}{|P|^2}.$

For many classical posets this is exactly solvable:

	chain C_N	$\binom{N+1}{2}$ ($\mathbb{P} \rightarrow \frac{1}{2}$)	
	Boolean B_N	3^N ($\mathbb{P} = (3/4)^N$)	
	Tamari $\mathcal{T}_N$	$\frac{2(4N+1)!}{(N+1)!(3N+2)!}$	[Chapoton '06]

A basic question about any finite poset

Given a poset $(P, \preceq)$, pick two elements independently and uniformly at random.

How likely are they comparable? $\mathbb{P}(X \preceq Y) = \frac{\#\{\text{intervals}\}}{|P|^2}.$

For many classical posets this is exactly solvable:

	chain C_N	$\binom{N+1}{2}$ ($\mathbb{P} \rightarrow \frac{1}{2}$)	
	Boolean B_N	3^N ($\mathbb{P} = (3/4)^N$)	
	Tamari $\mathcal{T}_N$	$\frac{2(4N+1)!}{(N+1)!(3N+2)!}$	[Chapoton '06]

For the **Weak Bruhat order**, write $\mathcal{W}_n = \{(\pi, \sigma) : \pi \preceq_W \sigma\}.$

A basic question about any finite poset

Given a poset $(P, \preceq)$, pick two elements independently and uniformly at random.

How likely are they comparable? $\mathbb{P}(X \preceq Y) = \frac{\#\{\text{intervals}\}}{|P|^2}.$

For many classical posets this is exactly solvable:

	chain C_N	$\binom{N+1}{2}$ ($\mathbb{P} \rightarrow \frac{1}{2}$)	
	Boolean B_N	3^N ($\mathbb{P} = (3/4)^N$)	
	Tamari $\mathcal{T}_N$	$\frac{2(4N+1)!}{(N+1)!(3N+2)!}$	[Chapoton '06]

For the **Weak Bruhat order**, write $\mathcal{W}_n = \{(\pi, \sigma) : \pi \preceq_W \sigma\}.$

Naive bounds

$$n! \leq |\mathcal{W}_n| \leq (n!)^2, \quad \text{i.e.} \\ (1+o(1)) n \log n \leq \log |\mathcal{W}_n| \leq (2+o(1)) n \log n.$$

The Big Question

$$(1 + o(1)) n \log n \leq \log |\mathcal{W}_n| \leq (2 + o(1)) n \log n$$

Hammett–Pittel (2008) [HP08] studied this and proved

$$(n!) \prod_{i=1}^n H(i) \leq |\mathcal{W}_n| \leq (n!)^2 (0.362)^n, \quad H(i) = \sum_{j \leq i} \frac{1}{j}.$$

Taking logarithms,

$$(1 + o(1)) n \log n + \log \log n \leq \log |\mathcal{W}_n| \leq (2 + o(1)) n \log n + \log(0.362) n.$$

Better than the naive bounds, but does not capture leading order behaviour.

The Big Question

$$(1 + o(1)) n \log n \leq \log |\mathcal{W}_n| \leq (2 + o(1)) n \log n$$

Hammett–Pittel (2008) [HP08] studied this and proved

$$(n!) \prod_{i=1}^n H(i) \leq |\mathcal{W}_n| \leq (n!)^2 (0.362)^n, \quad H(i) = \sum_{j \leq i} \frac{1}{j}.$$

Taking logarithms,

$$(1 + o(1)) n \log n + \log \log n \leq \log |\mathcal{W}_n| \leq (2 + o(1)) n \log n + \log(0.362) n.$$

Better than the naive bounds, but does not capture leading order behaviour.

A natural problem is to find the **coefficient of $n \log n$** : a constant $c_W \in [1, 2]$ with

$$\log |\mathcal{W}_n| = (c_W + o(1)) n \log n.$$

Main result

Theorem (B.–Cornejo–Hodges–Horn–Lesnevich–McAllister)

As $n \rightarrow \infty$,

$$|\mathcal{W}_n| = (n!)^2 \exp\left(-\left(\frac{1}{2} + o(1)\right) n \log n\right), \quad \mathbb{P}(\sigma_1 \preceq_W \sigma_2) = \exp\left(-\left(\frac{1}{2} + o(1)\right) n \log n\right).$$

Equivalently, c_W is determined exactly:

$$\log |\mathcal{W}_n| = \left(\frac{3}{2} + o(1)\right) n \log n \approx (n!)^{3/2}$$

$$\mathbb{P}(\sigma_1 \preceq_W \sigma_2) \approx 1/\sqrt{n!}$$

Main result

Theorem (B.–Cornejo–Hodges–Horn–Lesnevich–McAllister)

As $n \rightarrow \infty$,

$$|\mathcal{W}_n| = (n!)^2 \exp\left(-\left(\frac{1}{2} + o(1)\right) n \log n\right), \quad \mathbb{P}(\sigma_1 \preceq_W \sigma_2) = \exp\left(-\left(\frac{1}{2} + o(1)\right) n \log n\right).$$

Equivalently, c_W is determined exactly:

$$\log |\mathcal{W}_n| = \left(\frac{3}{2} + o(1)\right) n \log n \approx (n!)^{3/2}$$

$$\mathbb{P}(\sigma_1 \preceq_W \sigma_2) \approx 1/\sqrt{n!}$$

On the logarithmic scale, this is **exactly halfway** between the naive bounds $n!$ and $(n!)^2$.

Proof sketch, step 1: reduce to typical permutations


Count intervals by their top element:

$$|\mathcal{W}_n| = \sum_{\sigma \in \mathfrak{S}_n} e(\sigma), \quad e(\sigma) := \#\{\pi : \pi \preceq_W \sigma\}$$

is the size of the **down-set** of σ . We are left to compute:

how large is $e(\sigma)$ for a typical σ ?

down-set of $\sigma = 3421$ in $\mathfrak{S}_4$: $e(\sigma) = 12$



Proof sketch, step 1: reduce to typical permutations

Count intervals by their top element:

$$|\mathcal{W}_n| = \sum_{\sigma \in \mathfrak{S}_n} e(\sigma), \quad e(\sigma) := \#\{\pi : \pi \preceq_W \sigma\}$$

is the size of the **down-set** of σ . We are left to compute:


how large is $e(\sigma)$ for a typical σ ?

We will show: Typically,

$$e(\sigma) = (n!)^{1/2+o(1)} \approx \sqrt{n!},$$

hence $|\mathcal{W}_n| \approx n! \cdot \sqrt{n!} = (n!)^{3/2}$.

down-set of $\sigma = 3421$ in $\mathfrak{S}_4$: $e(\sigma) = 12$




Step 2: the poset $P(\sigma)$

From σ , build the poset $P(\sigma)$ on positions $\{1, \dots, n\}$:

$$i \prec j \iff i < j \text{ and } \sigma(i) < \sigma(j).$$

Its *chains* are exactly the **increasing subsequences** of σ . Comparable elements are **non-inversions**.



Hasse diagram of $P(2413)$.

Step 2: the poset $P(\sigma)$


From σ , build the poset $P(\sigma)$ on positions $\{1, \dots, n\}$:

$$i \prec j \iff i < j \text{ and } \sigma(i) < \sigma(j).$$

Its *chains* are exactly the **increasing subsequences** of σ . Comparable elements are **non-inversions**.

Key fact


Since weak order is containment of inversions, $e(\sigma)$ equals the number of *linear extensions* of $P(\sigma)$.



Hasse diagram of $P(2413)$.


Step 3: RSK and Greene's theorem

Greene's theorem [Greene '74]. For the RSK shape $\lambda(\sigma)$, the sum $\lambda_1 + \dots + \lambda_k$ is the largest total size of k disjoint chains of $P(\sigma)$ (increasing subsequences).



Step 3: RSK and Greene's theorem


Greene's theorem [Greene '74]. For the RSK shape $\lambda(\sigma)$, the sum $\lambda_1 + \dots + \lambda_k$ is the largest total size of k disjoint chains of $P(\sigma)$ (increasing subsequences).



Heuristic. If $P(\sigma)$ were a disjoint union of chains of lengths $\lambda_1, \dots, \lambda_r$, it would have $\frac{n!}{\lambda_1! \dots \lambda_r!}$ linear extensions; so $\log e(\sigma) \approx n \log n - \sum_i \lambda_i \log \lambda_i$. An appeal to the Bochkov–Petrov inequalities [Bochkov–Petrov '21] makes this precise.

Step 4: typical shape in Plancherel measure


For uniform σ , the shape $\lambda(\sigma)$ follows the **Plancherel measure** and concentrates on a fixed limit shape [Logan–Shepp '77; Vershik–Kerov '77]. Typically every part is $\lambda_i = \Theta(\sqrt{n})$ — no long rows or columns [Baik–Deift–Johansson '99].



Step 4: typical shape in Plancherel measure

For uniform σ , the shape $\lambda(\sigma)$ follows the **Plancherel measure** and concentrates on a fixed limit shape [Logan–Shepp '77; Vershik–Kerov '77]. Typically every part is $\lambda_i = \Theta(\sqrt{n})$ — no long rows or columns [Baik–Deift–Johansson '99]. So $\log \lambda_i \approx \frac{1}{2} \log n$, and since $\sum_i \lambda_i = n$,

$$\sum_i \lambda_i \log \lambda_i \approx \frac{1}{2} (\log n) \sum_i \lambda_i = \frac{1}{2} n \log n.$$




Step 4: typical shape in Plancherel measure

For uniform σ , the shape $\lambda(\sigma)$ follows the **Plancherel measure** and concentrates on a fixed limit shape [Logan–Shepp '77; Vershik–Kerov '77]. Typically every part is $\lambda_i = \Theta(\sqrt{n})$ — no long rows or columns [Baik–Deift–Johansson '99]. So $\log \lambda_i \approx \frac{1}{2} \log n$, and since $\sum_i \lambda_i = n$,

$$\sum_i \lambda_i \log \lambda_i \approx \frac{1}{2} (\log n) \sum_i \lambda_i = \frac{1}{2} n \log n.$$

Computing $\mathcal{W}_n$

$$\log e(\sigma) \approx n \log n - \sum_i \lambda_i \log \lambda_i \approx \frac{1}{2} n \log n \Rightarrow e(\sigma) \approx (n!)^{1/2},$$
$$|\mathcal{W}_n| \approx n! e(\sigma) \approx (n!)^{3/2}.$$




Strong Bruhat order

Again, write $\tau = s_{i_1} \cdots s_{i_k}$ as a *reduced* product of simple transpositions.

Definition (weak order)

$\sigma \preceq_W \tau$ iff σ is a **prefix** of some reduced word $\tau = s_{i_1} \cdots s_{i_k}$; that is,

$$\sigma = s_{i_1} \cdots s_{i_m}, \quad m \leq k$$




Strong Bruhat order

Again, write $\tau = s_{i_1} \cdots s_{i_k}$ as a *reduced* product of simple transpositions.

Definition (**weak** strong order)

$\sigma \preceq_W \tau$ iff σ is a **prefix** of some reduced word $\tau = s_{i_1} \cdots s_{i_k}$; that is,

$$\sigma = s_{i_1} \cdots s_{i_m}, \quad m \leq k$$




Strong Bruhat order

Again, write $\tau = s_{i_1} \cdots s_{i_k}$ as a *reduced* product of simple transpositions.

Definition (**weak** strong order)

$\sigma \preceq_W \preceq_B \tau$ iff σ is a **prefix subword** of some reduced word $\tau = s_{i_1} \cdots s_{i_k}$; that is,

$$\sigma = s_{i_1} \cdots s_{i_m}, \quad m \leq k$$




Strong Bruhat order

Again, write $\tau = s_{i_1} \cdots s_{i_k}$ as a *reduced* product of simple transpositions.

Definition (~~weak~~ strong order)

$\sigma \preceq_W \preceq_B \tau$ iff σ is a **prefix subword** of some reduced word $\tau = s_{i_1} \cdots s_{i_k}$; that is,

$$\begin{aligned} \sigma &= s_{i_1} \cdots s_{i_m}, \quad m \leq k \\ \sigma &= s_{i_{j_1}} \cdots s_{i_{j_m}}, \quad j_1 < \cdots < j_m \end{aligned}$$



Strong Bruhat order


Again, write $\tau = s_{i_1} \cdots s_{i_k}$ as a *reduced* product of simple transpositions.

Definition (**weak strong** order)

$\sigma \preceq_W \preceq_B \tau$ iff σ is a **prefix subword** of some reduced word $\tau = s_{i_1} \cdots s_{i_k}$; that is,

$$\begin{aligned} \sigma &= s_{i_1} \cdots s_{i_m}, \quad m \leq k \\ \sigma &= s_{i_{j_1}} \cdots s_{i_{j_m}}, \quad j_1 < \cdots < j_m \end{aligned}$$

- Weak refines strong: $\sigma \preceq_W \tau \implies \sigma \preceq_B \tau$.
- If $\sigma \preceq_B \tau$, then in fact *any* reduced expression for τ contains a subword for σ .
- Cover relations: an occurrence of the mesh pattern  becomes an occurrence of the mesh pattern .



Strong order and convergent research

Hammett–Pittel gave an *exponentially* decaying lower bound $(0.708)^n = \exp(\log(0.708)n)$; we improve it to a **subexponential** one:

Theorem (B.–Cornejo–Hodges–Horn–Lesnevich–McAllister)

$$\mathbb{P}(\sigma_1 \preceq_B \sigma_2) \geq \exp\left(- (6 + o(1))\sqrt{n}(\log n)^{3/2}\right) = \exp\left(- n^{1/2+o(1)}\right).$$

Strong order and convergent research

Hammett–Pittel gave an *exponentially* decaying lower bound $(0.708)^n = \exp(\log(0.708)n)$; we improve it to a **subexponential** one:

Theorem (B.–Cornejo–Hodges–Horn–Lesnevich–McAllister)

$$\mathbb{P}(\sigma_1 \preceq_B \sigma_2) \geq \exp\left(- (6 + o(1))\sqrt{n}(\log n)^{3/2}\right) = \exp\left(- n^{1/2+o(1)}\right).$$

Idea: model the *tableau criterion* by pairs of random walks on $\mathbb{Z}$, giving a large family of pairs comparable with probability $1 - o(1)$.

Strong order and convergent research

Hammett–Pittel gave an *exponentially* decaying lower bound $(0.708)^n = \exp(\log(0.708)n)$; we improve it to a **subexponential** one:

Theorem (B.–Cornejo–Hodges–Horn–Lesnevich–McAllister)

$$\mathbb{P}(\sigma_1 \preceq_B \sigma_2) \geq \exp\left(- (6 + o(1))\sqrt{n}(\log n)^{3/2}\right) = \exp\left(- n^{1/2+o(1)}\right).$$

Idea: model the *tableau criterion* by pairs of random walks on $\mathbb{Z}$, giving a large family of pairs comparable with probability $1 - o(1)$. *Just days later...*

Strong order and convergent research

Hammett–Pittel gave an *exponentially* decaying lower bound $(0.708)^n = \exp(\log(0.708)n)$; we improve it to a **subexponential** one:

Theorem (B.–Cornejo–Hodges–Horn–Lesnevich–McAllister)

$$\mathbb{P}(\sigma_1 \preceq_B \sigma_2) \geq \exp\left(- (6 + o(1))\sqrt{n}(\log n)^{3/2}\right) = \exp\left(- n^{1/2+o(1)}\right).$$

Idea: model the *tableau criterion* by pairs of random walks on $\mathbb{Z}$, giving a large family of pairs comparable with probability $1 - o(1)$. *Just days later...*

Theorem (Christo–Michelen)

$$\mathbb{P}(\sigma_1 \preceq_B \sigma_2) = \exp\left(- \Theta(\log^2 n)\right).$$

Also probabilistic, using a Gaussian process to model a *two-dimensional* random walk.


- Can we get asymptotics for other types?
- Can the asymptotics be refined by length of interval?
- What about more general chains?
- Is there a representation theory story here?

- Can we get asymptotics for other types?
- Can the asymptotics be refined by length of interval?
- What about more general chains?
- Is there a representation theory story here?

- Can we get asymptotics for other types?
- Can the asymptotics be refined by length of interval?
- What about more general chains?
- Is there a representation theory story here?

- Can we get asymptotics for other types?
- Can the asymptotics be refined by length of interval?
- What about more general chains?
- Is there a representation theory story here?

Thank you!



Comparability in Bruhat Orders — Boretsky, Cornejo, Hodges, Horn, Lesnevich, McAllister · [arXiv:2601.21158](https://arxiv.org/abs/2601.21158)

-  J. Boretsky, A. Cornejo, R. Hodges, P. Horn, N. Lesnevich, T. McAllister. *Comparability in Bruhat orders*. arXiv:2601.21158 (2026).
-  N. Christo, M. Michelen. *Comparability of random permutations in the strong Bruhat order*. arXiv:2602.16625 (2026).
-  A. Hammett, B. Pittel. *How often are two permutations comparable?* Trans. AMS **360** (2008).
-  I. Bochkov, F. Petrov. *Bounds for the number of linear extensions...* Order **38** (2021).
-  J. Baik, P. Deift, K. Johansson. *On the distribution of the length of the longest increasing subsequence...* JAMS **12** (1999).
-  C. Greene. *An extension of Schensted's theorem*. Adv. Math. **14** (1974).

-  F. Chapoton. *Sur le nombre d'intervalles dans les treillis de Tamari*. Sém. Lothar. Combin. (2006).
-  M. Bouvel, L. Ferrari, B. E. Tenner. *Between weak and Bruhat: the middle order on permutations*. Graphs Combin. **41** (2025).
-  A. Björner, F. Brenti. *Combinatorics of Coxeter groups*. GTM 231, Springer (2005).

A cautionary tale...

Thank you!



each letter: a copy of weak order on $\mathfrak{S}_5$, with several *intervals* (distinct colors) shaded in.

Comparability in Bruhat Orders — Boretsky, Cornejo, Hodges, Horn, Lesnevich, McAllister - arXiv:2601.21158

12/14