

Permutations containing a fixed number of occurrences of a pattern

Michael Waite
University of Florida

Permutation Patterns 2026
July 23, 2026

Basics

Definition

Define $S_{n,r}(q)$ to be the set of all permutations of length n that contain exactly r copies of q . Write $S_{n,0}(q)$ as $S_n(q)$.

Basics

Definition

Define $S_{n,r}(q)$ to be the set of all permutations of length n that contain exactly r copies of q . Write $S_{n,0}(q)$ as $S_n(q)$.

$p = 614235$

How many copies of 321?

Basics

Definition

Define $S_{n,r}(q)$ to be the set of all permutations of length n that contain exactly r copies of q . Write $S_{n,0}(q)$ as $S_n(q)$.

$p = 614235$

How many copies of 321?

Basics

Definition

Define $S_{n,r}(q)$ to be the set of all permutations of length n that contain exactly r copies of q . Write $S_{n,0}(q)$ as $S_n(q)$.

$p = 614235$

How many copies of 321?

Basics

Definition

Define $S_{n,r}(q)$ to be the set of all permutations of length n that contain exactly r copies of q . Write $S_{n,0}(q)$ as $S_n(q)$.

$p = 614235$



How many copies of 321?

2, so $p \in S_{6,2}(321)$.

Basics

Definition

Define $S_{n,r}(q)$ to be the set of all permutations of length n that contain exactly r copies of q . Write $S_{n,0}(q)$ as $S_n(q)$.

$p = 614235$



How many copies of 321?

2, so $p \in S_{6,2}(321)$.

How many copies of 231?

Basics

Definition

Define $S_{n,r}(q)$ to be the set of all permutations of length n that contain exactly r copies of q . Write $S_{n,0}(q)$ as $S_n(q)$.

$p = 614235$



How many copies of 321?

2, so $p \in S_{6,2}(321)$.

How many copies of 231?

0, so $p \in S_6(231)$.

Known Results

	$ S_{n,r}(321) $	$ S_{n,r}(231) $
$r = 1$	$\frac{3}{n} \binom{2n}{n-3}$ (Noonan 1996)	$\binom{2n-3}{n-3}$ (Bóna 1997)

Known Results

	$ S_{n,r}(321) $	$ S_{n,r}(231) $
$r = 1$	$\frac{3}{n} \binom{2n}{n-3}$ (Noonan 1996)	$\binom{2n-3}{n-3}$ (Bóna 1997)
$r = 2$	$\frac{59n^2+117n+100}{2n(n-1)(n+5)} \binom{2n}{n-4}$ (Fulmek 2001)	$\frac{n^3+17n^2-80n+80}{2n(n-1)} \binom{2n-6}{n-4}$ (Mansour, Vainshtein 2002)

Known Results

	$ S_{n,r}(321) $	$ S_{n,r}(231) $
$r = 1$	$\frac{3}{n} \binom{2n}{n-3}$ (Noonan 1996)	$\binom{2n-3}{n-3}$ (Bóna 1997)
$r = 2$	$\frac{59n^2+117n+100}{2n(n-1)(n+5)} \binom{2n}{n-4}$ (Fulmek 2001)	$\frac{n^3+17n^2-80n+80}{2n(n-1)} \binom{2n-6}{n-4}$ (Mansour, Vainshtein 2002)
$r = 3$	Enumeration known (Callan 2002)	Enumeration known (Mansour, Vainshtein 2002)
$r = 4$	Enumeration known (Callan 2002)	Enumeration known (Mansour, Vainshtein 2002)
$r = 5$		Enumeration known (Mansour, Vainshtein 2002)
$r = 6$		Enumeration known (Mansour, Vainshtein 2002)

Known Results

So we can do such enumerations for small r . But what can be said of general r ?

Known Results

So we can do such enumerations for small r . But what can be said of general r ?

Theorem (Bóna 1997)

Let $r \geq 1$. Let $S_r(z) = \sum_{n \geq 0} |S_{n,r}(231)| z^n$. Then

$$S_r(z) \in \mathbb{C}(z, \sqrt{1 - 4z}).$$

Known Results

So we can do such enumerations for small r . But what can be said of general r ?

Theorem (Bóna 1997)

Let $r \geq 1$. Let $S_r(z) = \sum_{n \geq 0} |S_{n,r}(231)|$. Then

$$S_r(z) \in \mathbb{C}(z, \sqrt{1 - 4z}).$$

For everything else, nothing so general is known about the generating functions.

Negative Results

What do we do if we try really hard to prove anything about a generating function, but nothing works?

Negative Results

What do we do if we try really hard to prove anything about a generating function, but nothing works?

Cry.

Negative Results

What do we do if we try really hard to prove anything about a generating function, but nothing works?

~~Cry.~~

Negative Results

What do we do if we try really hard to prove anything about a generating function, but nothing works?

~~Cry.~~

Maybe we can prove that the generating function is "bad".

Negative Results

What do we do if we try really hard to prove anything about a generating function, but nothing works?

~~Cry.~~

Maybe we can prove that the generating function is "bad". But how?

Negative Results

What do we do if we try really hard to prove anything about a generating function, but nothing works?

~~Cry.~~

Maybe we can prove that the generating function is "bad". But how?

Theorem

Let $f(z) = \sum_{n \geq 0} f_n z^n$ be a generating function. If α **is not a non-negative integer (is a negative integer)** and for all positive integers n the chain of inequalities

$$kn^\alpha \gamma^n \leq f_n \leq Kn^\alpha \gamma^n$$

holds, then $f(z)$ is **not a rational (is not an algebraic)** power series.

Negative Results

This means that if we can find bounds

$$k_r n^\alpha |S_n(q)| \leq |S_{n,r}(q)| \leq K_r n^\alpha |S_n(q)|$$

and if we know the asymptotics of $|S_n(q)|$, then we can make conclusions about the generating function for $|S_{n,r}(q)|$.

The Trivial Bound

Proposition

Let $r > 0$ and let q be any permutation. Then there are constants k_r and K_r so that


$$k_r |S_n(q)| \leq |S_{n,r}(q)| \leq K_r n^{2r} |S_n(q)|.$$

The Trivial Bound

Proposition

Let $r > 0$ and let q be any permutation. Then there are constants k_r and K_r so that

$$k_r |S_n(q)| \leq |S_{n,r}(q)| \leq K_r n^{2r} |S_n(q)|.$$




The Trivial Bound

Proposition

Let $r > 0$ and let q be any permutation. Then there are constants k_r and K_r so that

$$k_r |S_n(q)| \leq |S_{n,r}(q)| \leq K_r n^{2r} |S_n(q)|.$$





The Trivial Bound

Proposition

Let $r > 0$ and let q be any permutation. Then there are constants k_r and K_r so that

$$k_r |S_n(q)| \leq |S_{n,r}(q)| \leq K_r n^{2r} |S_n(q)|.$$





The Trivial Bound

Proposition

Let $r > 0$ and let q be any permutation. Then there are constants k_r and K_r so that

$$k_r |S_n(q)| \leq |S_{n,r}(q)| \leq K_r n^{2r} |S_n(q)|.$$



An Upper Bound for 321

Theorem (W. 2026+)

Let $r > 0$. Then there is a constant K_r so that

$$|S_{n,r}(321)| \leq K_r |S_n(321)|.$$

An Upper Bound for 321

Theorem (W. 2026+)

Let $r > 0$. Then there is a constant K_r so that

$$|S_{n,r}(321)| \leq K_r |S_n(321)|.$$

Definition

Define $S_{n,r}^q(321)$ to be the subset of $S_{n,r}(321)$ where the entries participating in 321 patterns form a q pattern.

An Upper Bound for 321

Theorem (W. 2026+)


Let $r > 0$. Then there is a constant K_r so that

$$|S_{n,r}(321)| \leq K_r |S_n(321)|.$$

Definition

Define $S_{n,r}^q(321)$ to be the subset of $S_{n,r}(321)$ where the entries participating in 321 patterns form a q pattern.

$p = 614235$



An Upper Bound for 321

Theorem (W. 2026+)


Let $r > 0$. Then there is a constant K_r so that

$$|S_{n,r}(321)| \leq K_r |S_n(321)|.$$

Definition

Define $S_{n,r}^q(321)$ to be the subset of $S_{n,r}(321)$ where the entries participating in 321 patterns form a q pattern.

$p = 614235$



An Upper Bound for 321

Theorem (W. 2026+)

Let $r > 0$. Then there is a constant K_r so that


$$|S_{n,r}(321)| \leq K_r |S_n(321)|.$$

Definition

Define $S_{n,r}^q(321)$ to be the subset of $S_{n,r}(321)$ where the entries participating in 321 patterns form a q pattern.

$p = 614235$

So $p \in S_{6,2}^{4312}(321)$.



An Upper Bound for 321


Now for each q , we will construct an injection

$$\phi^q : S_{n,r}^q(321) \hookrightarrow \bigcup_{k=1}^n S_{k-1+i,s}(321) \times S_{n-k+j,t}(321)$$

for some constants i, j and for some $s, t < r$.

An Upper Bound for 321

Now for each q , we will construct an injection


$$\phi^q : S_{n,r}^q(321) \hookrightarrow \bigcup_{k=1}^n S_{k-1+i,s}(321) \times S_{n-k+j,t}(321)$$


for some constants i, j and for some $s, t < r$.

Pairs of permutations on $n - 1 + i + j$ total elements,
with $s < r$ and $t < r$ copies of 321.

An Upper Bound for 321

Now for each q , we will construct an injection

$$\phi^q : S_{n,r}^q(321) \hookrightarrow \bigcup_{k=1}^n S_{k-1+i,s}(321) \times S_{n-k+j,t}(321)$$


for some constants i, j and for some $s, t < r$.


Pairs of permutations on $n - 1 + i + j$ total elements,
with $s < r$ and $t < r$ copies of 321.

By induction, the right hand side is at most $c_q |S_n(321)|$. So we get

$$|S_{n,r}^q(321)| \leq c_q |S_n(321)|.$$

An Upper Bound for 321

Now for each q , we will construct an injection

$$\phi^q : S_{n,r}^q(321) \hookrightarrow \bigcup_{k=1}^n S_{k-1+i,s}(321) \times S_{n-k+j,t}(321)$$


for some constants i, j and for some $s, t < r$.

Pairs of permutations on $n - 1 + i + j$ total elements,
with $s < r$ and $t < r$ copies of 321.


By induction, the right hand side is at most $c_q |S_n(321)|$. So we get

$$|S_{n,r}^q(321)| \leq c_q |S_n(321)|.$$


Since there are only finitely many possible q , we can sum over all possible q and get

$$|S_{n,r}(321)| \leq C_r |S_n(321)|.$$



The Map





The Map



The Map



The Map



A Generalization

Theorem (W 2026+)

Let q be a permutation, and let $r \geq 1$. Then there are constants k_r, K_r so that for all n sufficiently large we have

$$k_r |S_n(321 \ominus q)| \leq |S_{n,r}(321 \ominus q)| \leq K_r |S_n(321 \ominus q)|.$$

A Generalization

Theorem (W 2026+)

Let q be a permutation, and let $r \geq 1$. Then there are constants k_r, K_r so that for all n sufficiently large we have

$$k_r |S_n(321 \ominus q)| \leq |S_{n,r}(321 \ominus q)| \leq K_r |S_n(321 \ominus q)|.$$

Corollary (W 2026+)

Suppose $k \geq 3$. There is $c_{k,r}$ and $C_{k,r}$ such that

$$c_{k,r} \frac{(k-1)^{2n}}{n^{((k^2-2k)/2)}} \leq |S_{n,r}(k(k-1)\dots 1)| \leq C_{k,r} \frac{(k-1)^{2n}}{n^{((k^2-2k)/2)}}.$$

Let $S_r(z) = \sum_{n \geq 0} |S_{n,r}(q)| z^n$. Then:

- If k is odd, then $S_r(z)$ is not rational.
- If k is even, then $S_r(z)$ is not algebraic.

A Generalization

Theorem (W 2026+)

Let q be a permutation, and let $r \geq 1$. Then there are constants k_r, K_r so that for all n sufficiently large we have

$$k_r |S_n(321 \ominus q)| \leq |S_{n,r}(321 \ominus q)| \leq K_r |S_n(321 \ominus q)|.$$

Corollary (W 2026+)

Suppose $k \geq 3$. There is $c_{k,r}$ and $C_{k,r}$ such that

$$c_{k,r} \frac{(k-1)^{2n}}{n^{((k^2-2k)/2)}} \leq |S_{n,r}(k(k-1)\dots 1)| \leq C_{k,r} \frac{(k-1)^{2n}}{n^{((k^2-2k)/2)}}.$$

Let $S_r(z) = \sum_{n \geq 0} |S_{n,r}(q)| z^n$. Then:

- If k is odd, then $S_r(z)$ is not rational.
- If k is even, then $S_r(z)$ is not algebraic.

Bóna and Burstein (2021) proved this result for $r = 1$.

Further Results

Theorem (W 2026+)

Let q be a separable indecomposable permutation of length at least 2 which is of the form $q_1 \ominus \dots \ominus q_k$ for some $k \geq 2$ where all of the q_i are skew-indecomposable and $q_2, \dots, q_{k-1}$ are not 1. Then there are constants k and K such that for all n sufficiently large we have

$$kn^r |S_n(q)| \leq |S_{n,r}(q)| \leq Kn^r |S_n(q)|.$$

Further Results

Theorem (W 2026+)

Let q be a separable indecomposable permutation of length at least 2 which is of the form $q_1 \ominus \dots \ominus q_k$ for some $k \geq 2$ where all of the q_i are skew-indecomposable and $q_2, \dots, q_{k-1}$ are not 1. Then there are constants k and K such that for all n sufficiently large we have

$$kn^r |S_n(q)| \leq |S_{n,r}(q)| \leq Kn^r |S_n(q)|.$$

Corollary (W 2026+)

Suppose $r \geq 1$. Then there is c_k and C_k such that

$$c_k \frac{(k-1)^{2n}}{n^{((k^2-2k)/2)-r}} \leq |S_{n,r}(23\dots k1)| \leq C_k \frac{(k-1)^{2n}}{n^{((k^2-2k)/2)-r}}.$$

Let $S_r(z) = \sum_{n \geq 0} |S_{n,r}(23\dots k1)| z^n$. Then:

- If k is odd, then $S_r(z)$ is not rational.
- If k is even and $r < \frac{k^2-2k}{2}$, then $S_r(z)$ is not algebraic.

Further Results

Corollary (W 2026+)

Suppose $r \geq 1$. Then there is c and C such that

$$c \frac{9^n}{n^{4-r}} \leq |S_{n,r}(3412)| \leq C \frac{9^n}{n^{4-r}}.$$

Let $S(z) = \sum_{n \geq 0} |S_{n,r}(3412)| z^n$. If $r < 4$, then $S(z)$ is not algebraic.

Further Results

Corollary (W 2026+)

Suppose $r \geq 1$. Then there is c and C such that

$$c \frac{9^n}{n^{4-r}} \leq |S_{n,r}(3412)| \leq C \frac{9^n}{n^{4-r}}.$$

Let $S(z) = \sum_{n \geq 0} |S_{n,r}(3412)| z^n$. If $r < 4$, then $S(z)$ is not algebraic.

Corollary (W 2026+)

Suppose $r \geq 1$. Then there is c and C such that

$$c \frac{8^n}{n^{5/2-r}} \leq |S_{n,r}(4213)| \leq C \frac{8^n}{n^{5/2-r}}.$$

Let $S(z) = \sum_{n \geq 0} |S_{n,r}(4213)| z^n$. Then $S(z)$ is not rational.

Further Directions

Conjecture (Conway and Guttmann 2023)

Let $q \in \{4321, 4312\}$ and $r \geq 1$. Then there is a constant K so that

$$|S_{n,r}(q)| \sim K |S_n(q)|.$$

Let $q \in \{4123, 3412, 4231, 4213, 3142\}$ and $r \geq 1$. Then there is a constant K so that

$$|S_{n,r}(q)| \sim K n^r |S_n(q)|.$$

Further Directions

Conjecture (Conway and Guttmann 2023)

Let $q \in \{4321, 4312\}$ and $r \geq 1$. Then there is a constant K so that

$$|S_{n,r}(q)| \sim K |S_n(q)|.$$

Let $q \in \{4123, 3412, 4231, 4213, 3142\}$ and $r \geq 1$. Then there is a constant K so that

$$|S_{n,r}(q)| \sim K n^r |S_n(q)|.$$

Further Directions

Conjecture (Conway and Guttmann 2023)

Let $q \in \{4321, 4312\}$ and $r \geq 1$. Then there is a constant K so that

$$|S_{n,r}(q)| \sim K |S_n(q)|.$$

Let $q \in \{4123, 3412, 4231, 4213, 3142\}$ and $r \geq 1$. Then there is a constant K so that

$$|S_{n,r}(q)| \sim K n^r |S_n(q)|.$$

Further Directions

Conjecture (Conway and Guttmann 2023)

Let $q \in \{4321, 4312\}$ and $r \geq 1$. Then there is a constant K so that

$$|S_{n,r}(q)| \sim K |S_n(q)|.$$

Let $q \in \{4123, 3412, 4231, 4213, 3142\}$ and $r \geq 1$. Then there is a constant K so that

$$|S_{n,r}(q)| \sim K n^r |S_n(q)|.$$

Question

Is there a similar result for $|S_{n,r}(3142)|$ that generalizes well?

Further Directions

Conjecture (Conway and Guttmann 2023)

Let $q \in \{4321, 4312\}$ and $r \geq 1$. Then there is a constant K so that

$$|S_{n,r}(q)| \sim K |S_n(q)|.$$

Let $q \in \{4123, 3412, 4231, 4213, 3142\}$ and $r \geq 1$. Then there is a constant K so that

$$|S_{n,r}(q)| \sim K n^r |S_n(q)|.$$

Question

Is there a similar result for $|S_{n,r}(3142)|$ that generalizes well?

Further Directions

Conjecture (Conway and Guttmann 2023)

Let $q \in \{4321, 4312\}$ and $r \geq 1$. Then there is a constant K so that

$$|S_{n,r}(q)| \sim K |S_n(q)|.$$

Let $q \in \{4123, 3412, 4231, 4213, 3142\}$ and $r \geq 1$. Then there is a constant K so that

$$|S_{n,r}(q)| \sim K n^r |S_n(q)|.$$

Question

Is there a similar result for $|S_{n,r}(3142)|$ that generalizes well?

Question

What can be said about $|S_{n,r}(4312)|$?

Further Directions

In every case we have observed, either $|S_{n,r}(q)| = \Theta(|S_n(q)|)$ or $|S_{n,r}(q)| = \Theta(n^r |S_n(q)|)$. But it seems unlikely that this will always be true.

Further Directions

In every case we have observed, either $|S_{n,r}(q)| = \Theta(|S_n(q)|)$ or $|S_{n,r}(q)| = \Theta(n^r |S_n(q)|)$. But it seems unlikely that this will always be true.

Question

Is there a pattern q so that $|S_{n,r}(q)| = \Theta(n^{\alpha r} |S_n(q)|)$ for any $\alpha \notin \{0, 1\}$?



Thank you for listening!

Questions?