

Stack-Sorting Diagrams and Tableaux

Jerry Zhang

South Pasadena High School

July 20, 2026


West's Stack-Sorting Map

The stack-sorting map s is a deterministic function that sends permutations to permutations.

West's Stack-Sorting Map

The stack-sorting map s is a deterministic function that sends permutations to permutations.

Example:




$$s(32415) =$$

West's Stack-Sorting Map

The stack-sorting map s is a deterministic function that sends permutations to permutations.

Example:




$$s(32415) =$$

West's Stack-Sorting Map

The stack-sorting map s is a deterministic function that sends permutations to permutations.

Example:




$$s(32415) =$$

West's Stack-Sorting Map

The stack-sorting map s is a deterministic function that sends permutations to permutations.

Example:




$$s(32415) =$$

West's Stack-Sorting Map

The stack-sorting map s is a deterministic function that sends permutations to permutations.

Example:




$$s(32415) =$$

West's Stack-Sorting Map

The stack-sorting map s is a deterministic function that sends permutations to permutations.

Example:




$$s(32415) =$$

West's Stack-Sorting Map

The stack-sorting map s is a deterministic function that sends permutations to permutations.

Example:




$$s(32415) =$$

West's Stack-Sorting Map

The stack-sorting map s is a deterministic function that sends permutations to permutations.

Example:




$$s(32415) =$$

West's Stack-Sorting Map

The stack-sorting map s is a deterministic function that sends permutations to permutations.

Example:




$$s(32415) =$$

West's Stack-Sorting Map

The stack-sorting map s is a deterministic function that sends permutations to permutations.

Example:




$$s(32415) =$$

West's Stack-Sorting Map

The stack-sorting map s is a deterministic function that sends permutations to permutations.

Example:




$$s(32415) = 23145.$$

West's Stack-Sorting Map

The stack-sorting map s is a deterministic function that sends permutations to permutations.

Example:




$$s(32415) = 23145.$$

Try it for yourself: $s(314592687) = 134526789$.

West's Stack-Sorting Map

The stack-sorting map s is a deterministic function that sends permutations to permutations.

Example:



$$s(32415) = 23145.$$

Try it for yourself: $s(314592687) = 134526789$.

Observation. If a permutation π is of length n , then $s^{n-1}(\pi)$ is sorted.

We denote the minimum integer k for which $s^k(\pi)$ is sorted by $\langle \pi \rangle$, and a permutation π is *t-stack-sortable* if and only if $\langle \pi \rangle \leq t$.

Some Interesting Questions

Let $\mathcal{W}_t(n)$ denote the set of all t -stack-sortable permutations in S_n . What is $|\mathcal{W}_t(n)|$?

Some Interesting Questions

Let $\mathcal{W}_t(n)$ denote the set of all t -stack-sortable permutations in S_n . What is $|\mathcal{W}_t(n)|$?

t	$ \mathcal{W}_t(n) $	Reference
1	$C_n = \frac{1}{n+1} \binom{2n}{n}$	Knuth (1968)
2	$\frac{2}{(n+1)(2n+1)} \binom{3n}{n}$	Zeilberger (1992)
3	polynomial time	Defant (2020)
≥ 4	open	

Some Interesting Questions

Let $\mathcal{W}_t(n)$ denote the set of all t -stack-sortable permutations in S_n . What is $|\mathcal{W}_t(n)|$?

t	$ \mathcal{W}_t(n) $	Reference
1	$C_n = \frac{1}{n+1} \binom{2n}{n}$	Knuth (1968)
2	$\frac{2}{(n+1)(2n+1)} \binom{3n}{n}$	Zeilberger (1992)
3	polynomial time	Defant (2020)
≥ 4	open	

Theorem (West 1990)

A permutation is 1-stack-sortable if and only if it avoids the pattern 231. A permutation is 2-stack-sortable if and only if it avoids 2341 and 3241 patterns that are not part of a greater 35241 pattern.

Some Interesting Questions

For all $n \in \mathbb{N}$, let

$$\mathcal{D}_n := \frac{1}{n!} \sum_{\pi \in S_n} \langle \pi \rangle.$$

What is the asymptotic behavior of $\mathcal{D}_n/n$ as $n \rightarrow \infty$?

Some Interesting Questions

For all $n \in \mathbb{N}$, let

$$\mathcal{D}_n := \frac{1}{n!} \sum_{\pi \in S_n} \langle \pi \rangle.$$

What is the asymptotic behavior of $\mathcal{D}_n/n$ as $n \rightarrow \infty$?

Conjecture (West 1990)

The limit of $\mathcal{D}_n/n$ exists.

Some Interesting Questions

For all $n \in \mathbb{N}$, let

$$\mathcal{D}_n := \frac{1}{n!} \sum_{\pi \in S_n} \langle \pi \rangle.$$

What is the asymptotic behavior of $\mathcal{D}_n/n$ as $n \rightarrow \infty$?

Conjecture (West 1990)

The limit of $\mathcal{D}_n/n$ exists.

Theorem (Defant 2020)

$$0.62433 \approx \lambda \leq \liminf_{n \rightarrow \infty} \frac{\mathcal{D}_n}{n} \leq \limsup_{n \rightarrow \infty} \frac{\mathcal{D}_n}{n} \leq \frac{3}{5}(7 - 8 \ln 2) \approx 0.87289$$

Introducing S'_n

Let $S'_n := \{\rho 0 \mid \rho \in S_n\}$ and

$$\mathcal{D}'_n := \frac{1}{n!} \sum_{\rho \in S_n} \langle \rho 0 \rangle.$$

Introducing S'_n

Let $S'_n := \{\rho 0 \mid \rho \in S_n\}$ and

$$\mathcal{D}'_n := \frac{1}{n!} \sum_{\rho \in S_n} \langle \rho 0 \rangle.$$

If $0 \in \pi$, then let $C_\pi = (c_{\pi,j})_{j \geq 1}$ denote the sequence of right-to-left maxima of π to the left of 0. For all $\pi \in S'_n$, we write

$$\begin{aligned} s(\pi) &= s(b_{\pi,1} c_{\pi,1} b_{\pi,2} c_{\pi,2} \dots b_{\pi,|C_\pi|} c_{\pi,|C_\pi|} 0) \\ &= s(b_{\pi,1}) s(b_{\pi,2}) \dots s(b_{\pi,|C_\pi|}) \underbrace{0 c_{\pi,|C_\pi|} \dots c_{\pi,1}}_{\text{increasing tail}}. \end{aligned}$$

Introducing S'_n

Let $S'_n := \{\rho 0 \mid \rho \in S_n\}$ and

$$\mathcal{D}'_n := \frac{1}{n!} \sum_{\rho \in S_n} \langle \rho 0 \rangle.$$

If $0 \in \pi$, then let $C_\pi = (c_{\pi,j})_{j \geq 1}$ denote the sequence of right-to-left maxima of π to the left of 0. For all $\pi \in S'_n$, we write

$$\begin{aligned} s(\pi) &= s(b_{\pi,1} c_{\pi,1} b_{\pi,2} c_{\pi,2} \dots b_{\pi,|C_\pi|} c_{\pi,|C_\pi|} 0) \\ &= s(b_{\pi,1}) s(b_{\pi,2}) \dots s(b_{\pi,|C_\pi|}) \underbrace{0 c_{\pi,|C_\pi|} \dots c_{\pi,1}}_{\text{increasing tail}}. \end{aligned}$$

- 1 The set of elements added to the tail when s is applied to $s^{i-1}(\pi)$ is precisely $C_{s^{i-1}(\pi)}$.

Introducing S'_n

Let $S'_n := \{\rho 0 \mid \rho \in S_n\}$ and

$$\mathcal{D}'_n := \frac{1}{n!} \sum_{\rho \in S_n} \langle \rho 0 \rangle.$$

If $0 \in \pi$, then let $C_\pi = (c_{\pi,j})_{j \geq 1}$ denote the sequence of right-to-left maxima of π to the left of 0. For all $\pi \in S'_n$, we write

$$\begin{aligned} s(\pi) &= s(b_{\pi,1} c_{\pi,1} b_{\pi,2} c_{\pi,2} \dots b_{\pi,|C_\pi|} c_{\pi,|C_\pi|} 0) \\ &= s(b_{\pi,1}) s(b_{\pi,2}) \dots s(b_{\pi,|C_\pi|}) \underbrace{0 c_{\pi,|C_\pi|} \dots c_{\pi,1}}_{\text{increasing tail}}. \end{aligned}$$

- ① The set of elements added to the tail when s is applied to $s^{i-1}(\pi)$ is precisely $C_{s^{i-1}(\pi)}$.
- ② When every element of $[n]$ is added to the tail, π is sorted.

Introducing S'_n

Let $S'_n := \{\rho 0 \mid \rho \in S_n\}$ and

$$\mathcal{D}'_n := \frac{1}{n!} \sum_{\rho \in S_n} \langle \rho 0 \rangle.$$

If $0 \in \pi$, then let $C_\pi = (c_{\pi,j})_{j \geq 1}$ denote the sequence of right-to-left maxima of π to the left of 0. For all $\pi \in S'_n$, we write

$$\begin{aligned} s(\pi) &= s(b_{\pi,1} c_{\pi,1} b_{\pi,2} c_{\pi,2} \dots b_{\pi,|C_\pi|} c_{\pi,|C_\pi|} 0) \\ &= s(b_{\pi,1}) s(b_{\pi,2}) \dots s(b_{\pi,|C_\pi|}) \underbrace{0 c_{\pi,|C_\pi|} \dots c_{\pi,1}}_{\text{increasing tail}}. \end{aligned}$$

- ① The set of elements added to the tail when s is applied to $s^{i-1}(\pi)$ is precisely $C_{s^{i-1}(\pi)}$.
- ② When every element of $[n]$ is added to the tail, π is sorted.

Proposition (Defant 2020)

$$\lim_{n \rightarrow \infty} \left(\frac{\mathcal{D}_n}{n} - \frac{\mathcal{D}'_n}{n} \right) = 0$$

Example: $\pi = 9\ 3\ 10\ 7\ 8\ 2\ 6\ 1\ 4\ 5\ 0$

(9 3) 10 (7) 8 (2) 6 (1 4) 5 0

10
8
6
5

Example: $\pi = 9\ 3\ 10\ 7\ 8\ 2\ 6\ 1\ 4\ 5\ 0$

(9 3) 10 (7) 8 (2) 6 (1 4) 5 0

$\downarrow_s$

(3) 9 7 (2 1) 4 0 5 6 8 10

10	9
8	7
6	4
5	

Example: $\pi = 9\ 3\ 10\ 7\ 8\ 2\ 6\ 1\ 4\ 5\ 0$

(9 3) 10 (7) 8 (2) 6 (1 4) 5 0

$\downarrow_s$

(3) 9 7 (2 1) 4 0 5 6 8 10

$\downarrow_s$

3 (1) 2 0 4 5 6 7 8 9 10

10	9	3
8	7	2
6	4	
5		

Example: $\pi = 9\ 3\ 10\ 7\ 8\ 2\ 6\ 1\ 4\ 5\ 0$

(9 3) 10 (7) 8 (2) 6 (1 4) 5 0

$\downarrow_s$

(3) 9 7 (2 1) 4 0 5 6 8 10

$\downarrow_s$

3 (1) 2 0 4 5 6 7 8 9 10

$\downarrow_s$

1 0 2 3 4 5 6 7 8 9 10

10	9	3	1
8	7	2	
6	4		
5			

Example: $\pi = 9\ 3\ 10\ 7\ 8\ 2\ 6\ 1\ 4\ 5\ 0$

(9 3) 10 (7) 8 (2) 6 (1 4) 5 0

$\downarrow_s$

(3) 9 7 (2 1) 4 0 5 6 8 10

$\downarrow_s$

3 (1) 2 0 4 5 6 7 8 9 10

$\downarrow_s$

1 0 2 3 4 5 6 7 8 9 10

$\downarrow_s$

0 1 2 3 4 5 6 7 8 9 10

10	9	3	1
8	7	2	
6	4		
5			

Example: $\pi = 9\ 3\ 10\ 7\ 8\ 2\ 6\ 1\ 4\ 5\ 0$

(9 3) 10 (7) 8 (2) 6 (1 4) 5 0

$\downarrow_s$

(3) 9 7 (2 1) 4 0 5 6 8 10

$\downarrow_s$

3 (1) 2 0 4 5 6 7 8 9 10

$\downarrow_s$

1 0 2 3 4 5 6 7 8 9 10

$\downarrow_s$

0 1 2 3 4 5 6 7 8 9 10

10	9	3	1
8	7	2	
6	4		
5			

Note that there are $\langle \pi \rangle = 4$ columns.

Example: $\pi = 9\ 3\ 10\ 7\ 8\ 2\ 6\ 1\ 4\ 5\ 0$

(9 3) 10 (7) 8 (2) 6 (1 4) 5 0

$\downarrow_s$

(3) 9 7 (2 1) 4 0 5 6 8 10

$\downarrow_s$

3 (1) 2 0 4 5 6 7 8 9 10

$\downarrow_s$

1 0 2 3 4 5 6 7 8 9 10

$\downarrow_s$

0 1 2 3 4 5 6 7 8 9 10

10	9	3	1
8	7	2	
6	4		
5			

Note that there are $\langle \pi \rangle = 4$ columns.
How do we arrange the rows?

Towards the Construction of the Stack-Sorting Tableaux

Fix any permutation $\pi \in S'_n$. For all $i \in [n]$, let the *column* $\text{col}_\pi(i)$ be the pass on which i is added to the tail. Furthermore, if $\text{col}_\pi(i) = 1$, then let the *row* $\text{row}_\pi(i)$ be the index of i in $C_{\pi,1}$, and otherwise, let it be the same as the row of the first entry to the right of i in $s^{\text{col}_\pi(i)-2}(\pi)$ that is in column $\text{col}_\pi(i) - 1$.

Towards the Construction of the Stack-Sorting Tableaux

Fix any permutation $\pi \in S'_n$. For all $i \in [n]$, let the *column* $\text{col}_\pi(i)$ be the pass on which i is added to the tail. Furthermore, if $\text{col}_\pi(i) = 1$, then let the *row* $\text{row}_\pi(i)$ be the index of i in $C_{\pi,1}$, and otherwise, let it be the same as the row of the first entry to the right of i in $s^{\text{col}_\pi(i)-2}(\pi)$ that is in column $\text{col}_\pi(i) - 1$.

9 3 10 7 8 2 6 1 4 5 0

10
8
6
5

Towards the Construction of the Stack-Sorting Tableaux

Fix any permutation $\pi \in S'_n$. For all $i \in [n]$, let the *column* $\text{col}_\pi(i)$ be the pass on which i is added to the tail. Furthermore, if $\text{col}_\pi(i) = 1$, then let the *row* $\text{row}_\pi(i)$ be the index of i in $C_{\pi,1}$, and otherwise, let it be the same as the row of the first entry to the right of i in $s^{\text{col}_\pi(i)-2}(\pi)$ that is in column $\text{col}_\pi(i) - 1$.

9 3 10 7 8 2 6 1 4 5 0

$\downarrow_s$

3 9 7 2 1 4 0 5 6 8 10

10	9
8	7
6	
5	4

Towards the Construction of the Stack-Sorting Tableaux

Fix any permutation $\pi \in S'_n$. For all $i \in [n]$, let the *column* $\text{col}_\pi(i)$ be the pass on which i is added to the tail. Furthermore, if $\text{col}_\pi(i) = 1$, then let the *row* $\text{row}_\pi(i)$ be the index of i in $C_{\pi,1}$, and otherwise, let it be the same as the row of the first entry to the right of i in $s^{\text{col}_\pi(i)-2}(\pi)$ that is in column $\text{col}_\pi(i) - 1$.

9 3 10 7 8 2 6 1 4 5 0

$\downarrow_s$

3 9 7 2 1 4 0 5 6 8 10

$\downarrow_s$

3 1 2 0 4 5 6 7 8 9 10

10	9	3
8	7	
6		
5	4	2

Towards the Construction of the Stack-Sorting Tableaux

Fix any permutation $\pi \in S'_n$. For all $i \in [n]$, let the *column* $\text{col}_\pi(i)$ be the pass on which i is added to the tail. Furthermore, if $\text{col}_\pi(i) = 1$, then let the *row* $\text{row}_\pi(i)$ be the index of i in $C_{\pi,1}$, and otherwise, let it be the same as the row of the first entry to the right of i in $s^{\text{col}_\pi(i)-2}(\pi)$ that is in column $\text{col}_\pi(i) - 1$.

9 3 10 7 8 2 6 1 4 5 0

$\downarrow_s$

3 9 7 2 1 4 0 5 6 8 10

$\downarrow_s$

3 1 2 0 4 5 6 7 8 9 10

$\downarrow_s$

1 0 2 3 4 5 6 7 8 9 10

10	9	3	
8	7		
6			
5	4	2	1

Towards the Construction of the Stack-Sorting Tableaux

Fix any permutation $\pi \in S'_n$. For all $i \in [n]$, let the *column* $\text{col}_\pi(i)$ be the pass on which i is added to the tail. Furthermore, if $\text{col}_\pi(i) = 1$, then let the *row* $\text{row}_\pi(i)$ be the index of i in $C_{\pi,1}$, and otherwise, let it be the same as the row of the first entry to the right of i in $s^{\text{col}_\pi(i)-2}(\pi)$ that is in column $\text{col}_\pi(i) - 1$.

9 3 10 7 8 2 6 1 4 5 0

$\downarrow_s$

3 9 7 2 1 4 0 5 6 8 10

$\downarrow_s$

3 1 2 0 4 5 6 7 8 9 10

$\downarrow_s$

1 0 2 3 4 5 6 7 8 9 10

$\downarrow_s$

0 1 2 3 4 5 6 7 8 9 10

10	9	3	
8	7		
6			
5	4	2	1

Stack-Sorting Diagrams and Tableaux

For every permutation $\pi \in S'_n$, define the composition $\alpha_\pi \models n$ such that the j th part of α_π is the number of entries i satisfying $\text{row}_\pi(i) = j$. In our running example, $\alpha_\pi = (3, 2, 1, 4)$.

Stack-Sorting Diagrams and Tableaux

For every permutation $\pi \in S'_n$, define the composition $\alpha_\pi \models n$ such that the j th part of α_π is the number of entries i satisfying $\text{row}_\pi(i) = j$. In our running example, $\alpha_\pi = (3, 2, 1, 4)$.

For every composition α , define the *composition diagram*

$$D(\alpha) := \{(i, j) \in \mathbb{N}^2 : i \leq \alpha_j\}.$$

We impose a partial order relation $\geq_D$ on $D(\alpha)$ such that $(x, y) \geq_D (z, w)$ if and only if $x \leq z$ and $y \leq w$. We refer to $D(\alpha_\pi)$ as the *stack-sorting diagram* of π .

Stack-Sorting Diagrams and Tableaux

For every permutation $\pi \in S'_n$, define the composition $\alpha_\pi \models n$ such that the j th part of α_π is the number of entries i satisfying $\text{row}_\pi(i) = j$. In our running example, $\alpha_\pi = (3, 2, 1, 4)$.

For every composition α , define the *composition diagram*

$$D(\alpha) := \{(i, j) \in \mathbb{N}^2 : i \leq \alpha_j\}.$$

We impose a partial order relation $\geq_D$ on $D(\alpha)$ such that $(x, y) \geq_D (z, w)$ if and only if $x \leq z$ and $y \leq w$. We refer to $D(\alpha_\pi)$ as the *stack-sorting diagram* of π .

Now, define the function $T_\pi : [n] \rightarrow D(\alpha_\pi)$ such that

$$T_\pi(i) := (\text{col}_\pi(i), \text{row}_\pi(i))$$

for all $i \in [n]$. We refer to T_π as the *stack-sorting tableau* of π .

Stack-Sorting Diagrams and Tableaux


Theorem (Zhang 2026)

For every permutation $\pi \in S'_n$, the function T_π is a linear extension of the poset $(D(\alpha_\pi), \geq_D)$.

Stack-Sorting Diagrams and Tableaux

Theorem (Zhang 2026)

For every permutation $\pi \in S'_n$, the function T_π is a linear extension of the poset $(D(\alpha_\pi), \geq_D)$.




Hasse diagram of $D(3, 2, 1, 4)$

Stack-Sorting Diagrams and Tableaux

Theorem (Zhang 2026)

For every permutation $\pi \in S'_n$, the function T_π is a linear extension of the poset $(D(\alpha_\pi), \geq_D)$.



Hasse diagram of $D(3, 2, 1, 4)$

But how is this useful?

Hook-Length Formula for Stack-Sorting Tableaux

Example:

1	1	2	
1	1		
1			
1	1	3	6

Hook-Length Formula for Stack-Sorting Tableaux

Example: $h_\alpha(4, 4)$ for $\alpha = (3, 2, 1, 4)$

1	1	2	
1	1		
1			
1	1	3	6

Hook-Length Formula for Stack-Sorting Tableaux

Example: $h_{\alpha}(4, 4)$ for $\alpha = (3, 2, 1, 4)$

1	1	2	
1	1		
1			
1	1	3	6

Hook-Length Formula for Stack-Sorting Tableaux

Example: $h_\alpha(4, 4)$ for $\alpha = (3, 2, 1, 4)$

1	1	2	
1	1		
1			
1	1	3	6

Hook-Length Formula for Stack-Sorting Tableaux

Example: $h_{\alpha}(4, 4)$ for $\alpha = (3, 2, 1, 4)$

1	1	2	
1	1		
1			
1	1	3	6

Hook-Length Formula for Stack-Sorting Tableaux

Example: $h_\alpha(3, 1)$ for $\alpha = (3, 2, 1, 4)$

1	1	2	
1	1		
1			
1	1	3	6

Hook-Length Formula for Stack-Sorting Tableaux

Example: $h_\alpha(3, 1)$ for $\alpha = (3, 2, 1, 4)$

1	1	2	
1	1		
1			
1	1	3	6

Hook-Length Formula for Stack-Sorting Tableaux

Example: $h_\alpha(3, 1)$ for $\alpha = (3, 2, 1, 4)$

1	1	2	
1	1		
1			
1	1	3	6

Hook-Length Formula for Stack-Sorting Tableaux

Theorem (Zhang 2026)

For every linear extension T of a composition $\alpha \models n$, the number of permutations $\pi \in S'_n$ satisfying $T_\pi = T$ is

$$\prod_{(i,j) \in D(\alpha)} h_\alpha(i,j).$$

Hook-Length Formula for Stack-Sorting Tableaux

Theorem (Zhang 2026)

For every linear extension T of a composition $\alpha \models n$, the number of permutations $\pi \in S'_n$ satisfying $T_\pi = T$ is

$$\prod_{(i,j) \in D(\alpha)} h_\alpha(i,j).$$

Notice how the above formula solely depends on α , so the number of permutations $\pi \in S'_n$ satisfying $\alpha_\pi = \alpha$ is

$$\mathcal{L}(D(\alpha)) \cdot \prod_{(i,j) \in D(\alpha)} h_\alpha(i,j)$$

where $\mathcal{L}(D(\alpha))$ denotes the number of linear extensions of $D(\alpha)$.

Two Applications

Theorem (Zhang 2026)

Fix $t \in \mathbb{N}$. For all $n \in \mathbb{N}$, the number of t -stack-sortable permutations in S'_n is computable in $O(n^{3t+1})$ time.

Two Applications

Theorem (Zhang 2026)

Fix $t \in \mathbb{N}$. For all $n \in \mathbb{N}$, the number of t -stack-sortable permutations in S'_n is computable in $O(n^{3t+1})$ time.

Main idea: For every composition α , we can compute the number of permutations π such that $\alpha_\pi = \alpha$, and the stack-sorting depth of π is precisely the width of α . The rest follows from grouping compositions together based on shape statistics and doing clever dynamic programming.

Two Applications

Theorem (Zhang 2026)

The limit of $\mathcal{D}_n/n$ exists, and $(\mathcal{D}'_{n-1}/n)_{n=2}^\infty$ is a monotonically increasing sequence that converges to $\lim_{n \rightarrow \infty} \mathcal{D}_n/n$.

Two Applications

Theorem (Zhang 2026)

The limit of $\mathcal{D}_n/n$ exists, and $(\mathcal{D}'_{n-1}/n)_{n=2}^\infty$ is a monotonically increasing sequence that converges to $\lim_{n \rightarrow \infty} \mathcal{D}_n/n$.

Main idea: We can reformulate $\mathcal{D}'_n$ in the language of stack-sorting tableaux through algebraic manipulation.

Two Applications

Theorem (Zhang 2026)

The limit of $\mathcal{D}_n/n$ exists, and $(\mathcal{D}'_{n-1}/n)_{n=2}^\infty$ is a monotonically increasing sequence that converges to $\lim_{n \rightarrow \infty} \mathcal{D}_n/n$.

Main idea: We can reformulate $\mathcal{D}'_n$ in the language of stack-sorting tableaux through algebraic manipulation.

$$0.73286 \approx \frac{\mathcal{D}'_{45}}{46} \leq \lim_{n \rightarrow \infty} \frac{\mathcal{D}_n}{n} \leq \frac{3}{5}(7 - 8 \ln 2) \approx 0.87289.$$

Thank you!

Questions?

Jerry Zhang `jerrylezhang@gmail.com`

`arXiv:2604.10779, 2606.24110`