

A charge-induced DAG structure on permutations

Juan B. Gil

Cal State LA, July 2026

Penn State Altoona

jgil@psu.edu

Joint work (in progress) with L. Lapointe and M. Weiner

Charge and cocharge vectors

Let $\sigma \in S_n$ and let $i_k = \text{pos}_\sigma(k)$. Set

$$\mathbf{ch}(\sigma)_{i_1} = 0 = \overline{\mathbf{ch}}(\sigma)_{i_1}.$$

For $k = 2, 3, \dots, n$:

- If $i_k > i_{k-1}$ (value k is to the right of value $k-1$):

$$\mathbf{ch}(\sigma)_{i_k} = \mathbf{ch}(\sigma)_{i_{k-1}} + 1 \quad \text{and} \quad \overline{\mathbf{ch}}(\sigma)_{i_k} = \overline{\mathbf{ch}}(\sigma)_{i_{k-1}}.$$

- If $i_k < i_{k-1}$ (value k is to the left of value $k-1$):

$$\mathbf{ch}(\sigma)_{i_k} = \mathbf{ch}(\sigma)_{i_{k-1}} \quad \text{and} \quad \overline{\mathbf{ch}}(\sigma)_{i_k} = \overline{\mathbf{ch}}(\sigma)_{i_{k-1}} + 1.$$

Example $\sigma = 37158264$

σ	3	7	1	5	8	2	6	4
ch (σ)								

Example $\sigma = 37158264$

σ	3	7	1	5	8	2	6	4
ch (σ)			0					

Example $\sigma = 37158264$

σ	3	7	1	5	8	2	6	4
$\text{ch}(\sigma)$			0			1		

Example $\sigma = 37158264$

σ	3	7	1	5	8	2	6	4
$\text{ch}(\sigma)$	1		0			1		

Example $\sigma = 37158264$

σ	3	7	1	5	8	2	6	4
$\text{ch}(\sigma)$	1		0			1		2

Example $\sigma = 37158264$

σ	3	7	1	5	8	2	6	4
$\text{ch}(\sigma)$	1		0	2		1		2

Example $\sigma = 37158264$

σ	3	7	1	5	8	2	6	4
$\text{ch}(\sigma)$	1		0	2		1	3	2

Example $\sigma = 37158264$

σ	3	7	1	5	8	2	6	4
ch(σ)	1	3	0	2		1	3	2

Example $\sigma = 37158264$

σ	3	7	1	5	8	2	6	4
ch(σ)	1	3	0	2	4	1	3	2

Example $\sigma = 37158264$

σ	3	7	1	5	8	2	6	4
$\text{ch}(\sigma)$	1	3	0	2	4	1	3	2
$\overline{\text{ch}}(\sigma)$	1	3	0	2	3	0	2	1

Example $\sigma = 37158264$

σ	3	7	1	5	8	2	6	4
$\mathbf{ch}(\sigma)$	1	3	0	2	4	1	3	2
$\overline{\mathbf{ch}}(\sigma)$	1	3	0	2	3	0	2	1

$$\mathbf{ch}(\sigma) = (1, 3, 0, 2, 4, 1, 3, 2), \quad \overline{\mathbf{ch}}(\sigma) = (1, 3, 0, 2, 3, 0, 2, 1).$$

$$\mathbf{ch} + \overline{\mathbf{ch}} + \mathbf{1} = \sigma \quad \rightsquigarrow \quad \mathbf{ch}(\sigma) + \overline{\mathbf{ch}}(\sigma) = \binom{n}{2}$$

Example $\sigma = 37158264$

σ	3	7	1	5	8	2	6	4
$\mathbf{ch}(\sigma)$	1	3	0	2	4	1	3	2
$\overline{\mathbf{ch}}(\sigma)$	1	3	0	2	3	0	2	1

$$\mathbf{ch}(\sigma) = (1, 3, 0, 2, 4, 1, 3, 2), \quad \overline{\mathbf{ch}}(\sigma) = (1, 3, 0, 2, 3, 0, 2, 1).$$

$$\mathbf{ch} + \overline{\mathbf{ch}} + \mathbf{1} = \sigma \rightsquigarrow \mathbf{ch}(\sigma) + \overline{\mathbf{ch}}(\sigma) = \binom{n}{2}$$

In particular,

$$\mathbf{ch}(\mathrm{id}_n) = (0, 1, 2, \dots, n-1) \text{ and } \overline{\mathbf{ch}}(\mathrm{id}_n) = (0, 0, 0, \dots, 0).$$

Main properties

The map $\sigma \mapsto \mathbf{ch}(\sigma)$ is a bijection between S_n and the set of vectors $v = (v_1, v_2, \dots, v_n) \in \mathbb{N}_0^n$ satisfying:

1. The entries of v form a contiguous interval $\{0, 1, \dots, M\}$ for some $0 \leq M < n$.
2. For each $1 \leq k \leq M$, the leftmost $k - 1$ precedes the rightmost k .

- Interaction with symmetries:

$$\mathbf{ch}(\sigma) = \overline{\mathbf{ch}}(\sigma^r)^r, \quad \overline{\mathbf{ch}}(\sigma) + \mathbf{ch}(\sigma^c) = M \cdot \mathbf{1},$$

where $M = \max(\overline{\mathbf{ch}}(\sigma)) = \max(\mathbf{ch}(\sigma^c))$.

Other properties (related to pattern avoidance)

- $\max(\mathbf{ch}(\pi \oplus \tau)) = \max(\mathbf{ch}(\pi)) + \max(\mathbf{ch}(\tau)) + 1.$
- $\max(\mathbf{ch}(\pi \ominus \tau)) = \max(\mathbf{ch}(\pi)) + \max(\mathbf{ch}(\tau)).$
- If σ avoids any pattern in $\{132, 213, 231, 312\}$, then $\max(\mathbf{ch}(\sigma)) = \text{asc}(\sigma).$
- If σ avoids any pattern in $\{123, 321\}$, then $\max(\mathbf{ch}(\sigma)) = \text{asc}(\sigma^{-1}).$

Other properties (related to pattern avoidance)

- $\max(\mathbf{ch}(\pi \oplus \tau)) = \max(\mathbf{ch}(\pi)) + \max(\mathbf{ch}(\tau)) + 1.$
- $\max(\mathbf{ch}(\pi \ominus \tau)) = \max(\mathbf{ch}(\pi)) + \max(\mathbf{ch}(\tau)).$
- If σ avoids any pattern in $\{132, 213, 231, 312\}$, then $\max(\mathbf{ch}(\sigma)) = \text{asc}(\sigma).$
- If σ avoids any pattern in $\{123, 321\}$, then $\max(\mathbf{ch}(\sigma)) = \text{asc}(\sigma^{-1}).$
- For *separable* σ : $\max(\mathbf{ch}(\sigma)) = \text{asc}(\sigma).$

$\mathbf{ch}(2413) = (0, 1, 0, 1)$ and $\mathbf{ch}(3142) = (1, 0, 2, 1).$

Other properties (related to pattern avoidance)

- $\max(\mathbf{ch}(\pi \oplus \tau)) = \max(\mathbf{ch}(\pi)) + \max(\mathbf{ch}(\tau)) + 1.$
- $\max(\mathbf{ch}(\pi \ominus \tau)) = \max(\mathbf{ch}(\pi)) + \max(\mathbf{ch}(\tau)).$
- If σ avoids any pattern in $\{132, 213, 231, 312\}$, then $\max(\mathbf{ch}(\sigma)) = \text{asc}(\sigma).$
- If σ avoids any pattern in $\{123, 321\}$, then $\max(\mathbf{ch}(\sigma)) = \text{asc}(\sigma^{-1}).$
- For *separable* σ : $\max(\mathbf{ch}(\sigma)) = \text{asc}(\sigma).$
 $\mathbf{ch}(2413) = (0, 1, 0, 1)$ and $\mathbf{ch}(3142) = (1, 0, 2, 1).$
- $\sigma \in S_n$ has a nondecreasing charge vector if and only if σ is layered (avoids 231 and 312).

The charge-cocharge relation

For $\sigma, \tau \in S_n$, we write $\sigma \rightarrow \tau$ if the entries of the vector

$$w = \overline{\mathbf{ch}}(\sigma) + \mathbf{ch}(\tau)$$

are all distinct and $\max(w) - \min(w) = n - 1$.

w is a shifted permutation of $\{0, 1, \dots, n - 1\}$

The charge-cocharge relation

For $\sigma, \tau \in S_n$, we write $\sigma \rightarrow \tau$ if the entries of the vector

$$w = \overline{\mathbf{ch}}(\sigma) + \mathbf{ch}(\tau)$$

are all distinct and $\max(w) - \min(w) = n - 1$.

w is a shifted permutation of $\{0, 1, \dots, n-1\}$

- $\sigma \rightarrow \sigma$.
- If $m = \min(w)$, then $nm = \mathbf{ch}(\tau) - \mathbf{ch}(\sigma)$.

$$\sigma \rightarrow \tau \Rightarrow \mathbf{ch}(\tau) \equiv \mathbf{ch}(\sigma) \pmod{n}.$$

Symmetries and cyclage invariance

The relation is compatible with the standard symmetries:

$$\sigma \rightarrow \tau \implies \tau^r \rightarrow \sigma^r, \quad \tau^c \rightarrow \sigma^c, \quad \sigma^{rc} \rightarrow \tau^{rc}.$$

If $\sigma \rightarrow \tau$, then the permutation $\sigma \circ \tau^{-1}$ is nonseparable.

Symmetries and cyclage invariance

The relation is compatible with the standard symmetries:

$$\sigma \rightarrow \tau \implies \tau^r \rightarrow \sigma^r, \quad \tau^c \rightarrow \sigma^c, \quad \sigma^{rc} \rightarrow \tau^{rc}.$$

If $\sigma \rightarrow \tau$, then the permutation $\sigma \circ \tau^{-1}$ is nonseparable.

Cyclage invariance. If $\varrho = 23 \cdots n1$, then

$$\sigma \rightarrow \tau \implies \sigma \circ \varrho \rightarrow \tau \circ \varrho.$$

THEOREM

The relation $\rightarrow$ on S_n has no directed cycles other than loops:

$$\sigma_1 \rightarrow \sigma_2 \rightarrow \cdots \rightarrow \sigma_\ell \rightarrow \sigma_1 \implies \sigma_1 = \cdots = \sigma_\ell.$$

THEOREM

The relation $\rightarrow$ on S_n has no directed cycles other than loops:

$$\sigma_1 \rightarrow \sigma_2 \rightarrow \cdots \rightarrow \sigma_\ell \rightarrow \sigma_1 \implies \sigma_1 = \cdots = \sigma_\ell.$$

Sketch of Proof. Induction on n .

- By cyclage, assume $\sigma_1(n) = 1$.
- Summing the shifts m around the cycle forces every shift to vanish... which propagates $\sigma_i(n) = 1$ to all i .
- Delete the 1 to get a chain in S_{n-1} .

Proof for the case $\ell = 3$

Recall: For every edge $\alpha \rightarrow \beta$, $\overline{\mathbf{ch}}(\alpha) + \mathbf{ch}(\beta)$ is a permutation of $\{m, \dots, m+n-1\}$, where $nm = \mathbf{ch}(\beta) - \mathbf{ch}(\alpha) \geq 0$.

Let $a, b, c \geq 0$ be the shifts of $\sigma \rightarrow \tau$, $\tau \rightarrow \pi$ and $\pi \rightarrow \sigma$. Then

$$n(a + b + c) = (\mathbf{ch}(\tau) - \mathbf{ch}(\sigma)) + (\mathbf{ch}(\pi) - \mathbf{ch}(\tau)) + (\mathbf{ch}(\sigma) - \mathbf{ch}(\pi)) = 0$$

so $a = b = c = 0$. Each of the three vectors is a permutation of $\{0, 1, \dots, n-1\}$; in particular each has *exactly one zero* entry.

By cyclage invariance we may assume $\sigma(n) = 1$, so

$$\overline{\mathbf{ch}}(\sigma)_n = 0, \quad \overline{\mathbf{ch}}(\sigma)_j \geq 1 \quad (j < n).$$

Case $\ell = 3$ continued

Since the vector $w = \overline{\mathbf{ch}}(\sigma) + \mathbf{ch}(\tau)$ has a unique zero entry, we must have $\mathbf{ch}(\tau)_n = 0$, and so $\tau(n) = 1$.

Applying the same argument to the edge $\tau \rightarrow \pi$ gives $\pi(n) = 1$.

All three permutations have the 1 in position n . Deleting it and standardizing, yields a cycle

$$\sigma' \rightarrow \tau' \rightarrow \pi' \rightarrow \sigma' \quad \text{in } S_{n-1}.$$

By the induction hypothesis $\sigma' = \tau' = \pi'$, and reinserting the 1 in the last position gives $\sigma = \tau = \pi$. □

Interlude: where the question came from

Luc needed, inside the **affine 0-Hecke algebra** $H^{(n)}$ (over $\mathbb{C}[x_1^\pm, \dots, x_n^\pm]$), a *complete family of orthogonal idempotents*.

His building block attaches to each σ the element

$$\psi_\sigma = (-1)^{\ell(\sigma) + \ell(w_0)} x^{\mathbf{ch}(\sigma)} \partial_{w_0} x^{\overline{\mathbf{ch}}(\sigma)}, \quad \partial_{w_0} = \frac{1}{a_\delta} \sum_w (-1)^{\ell(w)} w.$$

The charge and cocharge appear as **exponent vectors** of Laurent monomials. A single number wouldn't do — vectors are needed

Our combinatorics vs. his algebra

The core computation is $\partial_{w_0} x^\mu \partial_{w_0} = \frac{a_\mu}{a_\delta} \partial_{w_0}$, which gives

$$\psi_w \psi_\sigma \propto \frac{a_{\overline{\mathbf{ch}}(w) + \mathbf{ch}(\sigma)}}{a_\delta}, \quad \text{and} \quad a_\mu = 0 \iff \mu \text{ has a repeated entry.}$$

$$\mathbf{ch} + \overline{\mathbf{ch}} + \mathbf{1} = \sigma \iff \psi_\sigma^2 = \psi_\sigma \text{ (idempotency)}$$

$$\overline{\mathbf{ch}}(w) + \mathbf{ch}(\sigma) \text{ distinct} \iff \psi_w \psi_\sigma \neq 0 \text{ } (w \rightarrow \sigma)$$

$$\text{acyclicity of the arrow graph} \iff \text{chain products vanish}$$

Acyclicity makes the sum

$$L_\sigma = \sum_{\text{chains}} (-1)^\ell \psi_\sigma \psi_{w(1)} \cdots \psi_{w(\ell)} \text{ finite, and}$$

$\{L_w\}$ is a complete set of orthogonal idempotents with $\sum_w L_w = 1$.

Charge classes and the DAG

Total charge partitions S_n into n residue classes mod n , each of size $(n-1)!$.


Within each residue class, $\sigma \rightarrow \tau$ ($\sigma \neq \tau$) defines a **directed acyclic graph** on $(n-1)!$ vertices.

Charge classes and the DAG

Total charge partitions S_n into n residue classes mod n , each of size $(n-1)!$.

Within each residue class, $\sigma \rightarrow \tau$ ($\sigma \neq \tau$) defines a **directed acyclic graph** on $(n-1)!$ vertices.

Example: $n = 4$, class $ch \equiv 0 \pmod{4}$



Graph for $n = 5$, $ch \equiv 0 \pmod{5}$


$ch = 10$

$ch = 5$

$ch = 0$



Graph for $n = 6$



A new poset $\mathcal{P} = (S_n, \preceq)$

Call i a *long ascent* of w if $w(i+1) > w(i) + 1$.


Covering relations: $w \succ w s_i$ at each long ascent i ($1 \leq i \leq n-1$), together with the **affine cover** $w \succ \sigma_0 w$ when $w(1) > w(n) + 1$, where σ_0 swaps the two end positions.

$\mathcal{P} = (S_n, \preceq)$ is **ranked** with exactly n connected components: the level sets of total charge mod n . Each has $(n-1)!$ elements and a **unique minimum and maximum**.

New structures ($n = 4$, $\text{ch} \equiv 0$)

the poset $\mathcal{P}$

rk 0



rk 4

$n = 5$, $ch \equiv 0$

$rk = 0$

$rk = 1$

$rk = 2$

$rk = 3$

$rk = 4$

$rk = 5$


$rk = 6$

$rk = 7$

$rk = 8$

$rk = 9$

$rk = 10$



Why $\mathcal{P}$ is the right object

Read one-line notations *cyclically*: deleting $1, \dots, k$ from w leaves a cyclic word w_k on $\{k+1, \dots, m\}$. Then

$$rk(w) = \sum_{k=1}^{m-2} \text{ch}(w_k), \quad rk(w_0) = 0.$$

(The rank is *not* the total charge!)

Why $\mathcal{P}$ is the right object


Read one-line notations *cyclically*: deleting $1, \dots, k$ from w leaves a cyclic word w_k on $\{k+1, \dots, m\}$. Then

$$rk(w) = \sum_{k=1}^{m-2} \text{ch}(w_k), \quad rk(w_0) = 0.$$


(The rank is *not* the total charge!)

Conjecture: The rank strictly **increases along every arrow** of the charge–cocharge graph.


Graph for $n = 6$



Charge-cocharge arrows for $n = 6$



Charge-cocharge arrows for $n = 6$



References

- A. Lascoux, M.-P. Schützenberger, *Sur une conjecture de H. O. Foulkes*, C. R. Acad. Sci. Paris **286** (1978), A323–A324.
- L. Lapointe, *A complete family of idempotents in the affine 0-Hecke algebra*, preprint.

Thank you!