

On the Schur positivity of small descent enumerators for pattern avoiding sets

Kyle Celano


Wake Forest University

Permutation Patterns 4!

Descents and Small Descents

Let $\mathfrak{S}_n := \{\text{permutations } \sigma \text{ of } [n] = \{1, 2, \dots, n\}\}$.

A **small descent** of σ is a position $i \in [n-1]$ such that $\sigma_i = \sigma_{i+1} + 1$. Let $\text{SDES}(\sigma)$ be the set of small descents.




$$\text{SDES}(\sigma) = \{2, 5, 6\}$$

Descents and Small Descents

Let $\mathfrak{S}_n := \{\text{permutations } \sigma \text{ of } [n] = \{1, 2, \dots, n\}\}$.

A **small descent** of σ is a position $i \in [n-1]$ such that $\sigma_i = \sigma_{i+1} + 1$. Let $\text{SDES}(\sigma)$ be the set of small descents.



$$\text{SDES}(\sigma) = \{2, 5, 6\}$$

Small descents

Set $\text{sdes}(\sigma) := |\text{SDES}(\sigma)|$

Ex. $n = 4$.

k	$\text{sdes}(\pi) = k$
0	1234, 1342, 1423, 2314, 2341, 2413, 3124, 3142, 3412, 4123, 4231
1	1243, 1324, 2134, 2431, 3241, 3421, 4132, 4213, 4312
2	1432, 2143, 3214
3	4321

Small descents

Ex. $n = 4$. Let's look at $\text{SDES}(\pi) \subseteq [n - 1]$ directly

S	$\text{SDES}(\pi) = S$
$\emptyset$	1234, 1342, 1423, 2314, 2341, 2413, 3124, 3142, 3412, 4123, 4231
$\{1\}$	2134, 3241, 4312
$\{2\}$	1324, 2431, 4213
$\{3\}$	1243, 3421, 4132
$\{1, 2\}$	3214
$\{2, 3\}$	2143
$\{1, 3\}$	1432
$\{1, 2, 3\}$	4321

Small descents

Ex. $n = 4$. Let's look at $\text{SDES}(\pi) \subseteq [n - 1]$ directly

S	$\text{SDES}(\pi) = S$
$\emptyset$	1234, 1342, 1423, 2314, 2341, 2413, 3124, 3142, 3412, 4123, 4231
$\{1\}$	2134, 3241, 4312
$\{2\}$	1324, 2431, 4213
$\{3\}$	1243, 3421, 4132
$\{1, 2\}$	3214
$\{2, 3\}$	2143
$\{1, 3\}$	1432
$\{1, 2, 3\}$	4321

$|\{\pi \in \mathfrak{S}_n : \text{SDES}(\pi) = S\}|$ seems to only depend on $|S|$

Small descents

Ex. $n = 4$. Let's look at $\text{SDES}(\pi) \subseteq [n - 1]$ directly

S	$\text{SDES}(\pi) = S$
$\emptyset$	1234, 1342, 1423, 2314, 2341, 2413, 3124, 3142, 3412, 4123, 4231
$\{1\}$	2134, 3241, 4312
$\{2\}$	1324, 2431, 4213
$\{3\}$	1243, 3421, 4132
$\{1, 2\}$	3214
$\{2, 3\}$	2143
$\{1, 3\}$	1432
$\{1, 2, 3\}$	4321

Proposition

$|\{\pi \in \mathfrak{S}_n : \text{SDES}(\pi) = S\}|$ *only depends on* $|S|$.

Small descents

Ex. $n = 4$. Consider $\text{Av}_4(312)$

S	$\text{SDES}(\pi) = S$
$\emptyset$	1234, 1342, 1423 , 2314, 2341, 2413 , 3124 , 3142 , 3412 , 4123 , 4231
$\{1\}$	2134, 3241, 4312
$\{2\}$	1324, 2431, 4213
$\{3\}$	1243, 3421, 4132
$\{1, 2\}$	3214
$\{2, 3\}$	2143
$\{1, 3\}$	1432
$\{1, 2, 3\}$	4321

Small descents

Ex. $n = 4$. Consider $\text{Av}_4(312)$

S	$\text{SDES}(\pi) = S$
$\emptyset$	1234, 1342, 2314, 2341
$\{1\}$	2134, 3241
$\{2\}$	1324, 2431
$\{3\}$	1243, 3421
$\{1, 2\}$	3214
$\{2, 3\}$	2143
$\{1, 3\}$	1432
$\{1, 2, 3\}$	4321

Small descents

Ex. $n = 4$. Consider $\text{Av}_4(312)$

S	$\text{SDES}(\pi) = S$
$\emptyset$	1234, 1342, 2314, 2341
$\{1\}$	2134, 3241
$\{2\}$	1324, 2431
$\{3\}$	1243, 3421
$\{1, 2\}$	3214
$\{2, 3\}$	2143
$\{1, 3\}$	1432
$\{1, 2, 3\}$	4321

Proposition

$|\{\pi \in \text{Av}_n(312) : \text{SDES}(\pi) = S\}|$ *only depends on* $|S|$.

Small descents

Ex. $n = 4$. Consider $\text{Av}_4(312, 123)$

S	$\text{SDES}(\pi) = S$
$\emptyset$	$\emptyset$
$\{1\}$	3241
$\{2\}$	2431
$\{3\}$	3421
$\{1, 2\}$	3214
$\{2, 3\}$	2143
$\{1, 3\}$	1432
$\{1, 2, 3\}$	4321

Proposition

$|\{\pi \in \text{Av}_n(312, 123) : \text{SDES}(\pi) = S\}|$ *only depends on* $|S|$.


Theorem (C. +26)

Let Π be a finite set of patterns such that $\text{SDES}(\pi) = \emptyset$ for all $\pi \in \Pi$. Then, for all $S \subseteq [n-1]$, $|\{\pi \in \text{Av}_n(\Pi) : \text{SDES}(\pi) = S\}|$ only depends on $|S|$.


I.E. Any two sets of the same size are equally likely to be small descent sets

Proof


For a given $S \subseteq [n-1]$, one shows a bijection
 $\{\sigma \in \text{Av}_n(\Pi) : \text{SDES}(\sigma) = S\} \leftrightarrow \{u \in \text{Av}_{n-|S|}(\Pi) : \text{SDES}(u) = \emptyset\}$



(a) σ



(b) u (spread out)



(c) u

Observe: σ contains small-descent-avoiding pattern iff u does. □

Theorem (C. +26)

Let Π be a finite set of patterns such that $\text{SDES}(\pi) = \emptyset$ for all $\pi \in \Pi$. Then, for all $S \subseteq [n-1]$, $|\{\pi \in \text{Av}_n(\Pi) : \text{SDES}(\pi) = S\}|$ only depends on $|S|$.

Theorem (C. +26)

Let Π be a finite set of patterns such that $\text{SDES}(\pi) = \emptyset$ for all $\pi \in \Pi$. Then, for all $S \subseteq [n-1]$, $|\{\pi \in \text{Av}_n(\Pi) : \text{SDES}(\pi) = S\}|$ only depends on $|S|$.

Q: Why is this an “interesting” property?

Theorem (C. +26)

Let Π be a finite set of patterns such that $\text{SDES}(\pi) = \emptyset$ for all $\pi \in \Pi$. Then, for all $S \subseteq [n-1]$, $|\{\pi \in \text{Av}_n(\Pi) : \text{SDES}(\pi) = S\}|$ only depends on $|S|$.

Q: Why is this an “interesting” property?

A: Representation theory.

Theorem (C. +26)

Let Π be a finite set of patterns such that $\text{SDES}(\pi) = \emptyset$ for all $\pi \in \Pi$. Then, for all $S \subseteq [n-1]$, $|\{\pi \in \text{Av}_n(\Pi) : \text{SDES}(\pi) = S\}|$ only depends on $|S|$.

Q: Why is this an “interesting” property?

A: Representation theory.

Punchline: This theorem tells you that there is an action of the symmetric group on $\text{Av}_n(\Pi)$ that “decomposes” via small descents.

Symmetric Functions

Symmetric function: a (homogenous) polynomial f in $x_1, x_2, x_3, \dots$ that is invariant under any permutation $\sigma : \mathbb{Z}_{>0} \rightarrow \mathbb{Z}_{>0}$ of its indices i.e. $f(x_1, x_2, x_3, \dots) = f(x_{\sigma(1)}, x_{\sigma(2)}, x_{\sigma(3)}, \dots)$

$$2x_1^2x_2 + 2x_1x_2^2 + 2x_1^3x_3 + 2x_1x_3^2 + \dots \quad \checkmark$$

$$2x_1^2x_2 - 3x_1x_2^2 + 2x_1^3x_3 + 2x_1x_3^2 + \dots \quad \times$$

Schur functions

Semistandard Young tableaux (SSYT) of shape λ : a filling of a diagram of λ by positive integers that has weakly increasing rows and strictly increasing columns.

1	2	2	8	8
2	3	9	9	21
6	21			
21				

(a) semistandard Young tableau T

$$x^T := \prod_{i \geq 1} x_i^{\# \text{ of } i \text{ in } T}$$

$$x_1 x_2^3 x_3 x_6 x_8^2 x_9^2 x_{21}^3$$

(b) the monomial x^T

Schur function s_λ of shape λ

$$s_\lambda = \sum_{T \in \text{SSYT}_\lambda} x^T$$

Schur functions and representation theory

Schur functions correspond to the irreducible representations of the symmetric group.

A nonnegative integer linear combinations of Schur functions (a.k.a. **Schur positive** function) correspond to representations of the symmetric group.

Fundamental quasisymmetric functions

For a set $S = \{s_1 < s_2 < \cdots < s_{k-1}\} \subseteq [n-1]$,

$$F_{n,S} = \sum_{\substack{i_1 \leq \cdots \leq i_n \\ i_j < i_{j+1} \forall j \in S}} x_{i_1} \cdots x_{i_n}$$

$$F_{4,\{1\}} = x_1 x_2 x_2 x_2 + x_1 x_2 x_2 x_3 + x_1 x_2 x_3 x_3 + x_1 x_2 x_3 x_4 + \cdots$$

$$F_{4,\{2\}} = x_1 x_1 x_2 x_2 + x_1 x_2 x_2 x_2 + x_1 x_1 x_2 x_3 + x_1 x_2 x_3 x_4 + \cdots$$

$$F_{4,\{3\}} = x_1 x_1 x_1 x_2 + x_1 x_2 x_2 x_3 + x_1 x_1 x_2 x_3 + x_1 x_2 x_3 x_4 + \cdots$$

Fundamental quasisymmetric functions

For a set $S = \{s_1 < s_2 < \cdots < s_{k-1}\} \subseteq [n-1]$,

$$F_{n,S} = \sum_{\substack{i_1 \leq \cdots \leq i_n \\ i_j < i_{j+1} \forall j \in S}} x_{i_1} \cdots x_{i_n}$$

$$F_{4,\{1\}} = x_1 x_2 x_2 x_2 + x_1 x_2 x_2 x_3 + x_1 x_2 x_3 x_3 + x_1 x_2 x_3 x_4 + \cdots$$

$$F_{4,\{2\}} = x_1 x_1 x_2 x_2 + x_1 x_2 x_2 x_2 + x_1 x_1 x_2 x_3 + x_1 x_2 x_3 x_4 + \cdots$$

$$F_{4,\{3\}} = x_1 x_1 x_1 x_2 + x_1 x_2 x_2 x_3 + x_1 x_1 x_2 x_3 + x_1 x_2 x_3 x_4 + \cdots$$

Proposition (Gessel)

The quasisymmetric function

$$Q_{\mathfrak{S}_n}(\mathbf{x}) = \sum_{\sigma \in \mathfrak{S}_n} F_{n, \text{DES}(\sigma)}$$

is symmetric and moreover Schur positive.

Fundamental quasisymmetric functions

For a set $S = \{s_1 < s_2 < \cdots < s_{k-1}\} \subseteq [n-1]$,

$$F_{n,S} = \sum_{\substack{i_1 \leq \cdots \leq i_n \\ i_j < i_{j+1} \forall j \in S}} x_{i_1} \cdots x_{i_n}$$

$$F_{4,\{1\}} = x_1 x_2 x_2 x_2 + x_1 x_2 x_2 x_3 + x_1 x_2 x_3 x_3 + x_1 x_2 x_3 x_4 + \cdots$$

$$F_{4,\{2\}} = x_1 x_1 x_2 x_2 + x_1 x_2 x_2 x_2 + x_1 x_1 x_2 x_3 + x_1 x_2 x_3 x_4 + \cdots$$

$$F_{4,\{3\}} = x_1 x_1 x_1 x_2 + x_1 x_2 x_2 x_3 + x_1 x_1 x_2 x_3 + x_1 x_2 x_3 x_4 + \cdots$$

Proposition (Gessel)

The quasisymmetric function

$$Q_{\mathfrak{S}_n}(\mathbf{x}) = \sum_{\sigma \in \mathfrak{S}_n} F_{n, \text{DES}(\sigma)} = \sum_{\lambda \vdash n} |\text{SYT}_\lambda| s_\lambda = (x_1 + x_2 + \cdots)^n$$

is symmetric and moreover Schur positive. $Q_{\mathfrak{S}_n}(\mathbf{x})$ corresponds to the action of left multiplication of $\mathfrak{S}_n$ on itself.

Schur positive (classical) descent enumerators

For a subset $A \subseteq \mathfrak{S}_n$, define

$$\mathcal{Q}_A(\mathbf{x}) = \sum_{\pi \in A} F_{n, \text{DES}(\sigma)}$$

What subsets of permutations produce symmetric functions?
Which are Schur positive?

Schur positive (classical) descent enumerators

For a subset $A \subseteq \mathfrak{S}_n$, define

$$\mathcal{Q}_A(\mathbf{x}) = \sum_{\pi \in A} F_{n, \text{DES}(\sigma)}$$

What subsets of permutations produce symmetric functions?

Which are Schur positive?

- Gessel: With fixed inverse descent set

Schur positive (classical) descent enumerators

For a subset $A \subseteq \mathfrak{S}_n$, define

$$\mathcal{Q}_A(\mathbf{x}) = \sum_{\pi \in A} F_{n, \text{DES}(\sigma)}$$

What subsets of permutations produce symmetric functions?

Which are Schur positive?

- Gessel: With fixed inverse descent set
- Gessel–Reutenaur: With fixed cycle type

Schur positive (classical) descent enumerators

For a subset $A \subseteq \mathfrak{S}_n$, define

$$\mathcal{Q}_A(\mathbf{x}) = \sum_{\pi \in A} F_{n, \text{DES}(\sigma)}$$

What subsets of permutations produce symmetric functions?

Which are Schur positive?

- Gessel: With fixed inverse descent set
- Gessel–Reutenaur: With fixed cycle type
- Elizalde–Roichman: Arc permutations

Schur positive (classical) descent enumerators

For a subset $A \subseteq \mathfrak{S}_n$, define

$$\mathcal{Q}_A(\mathbf{x}) = \sum_{\pi \in A} F_{n, \text{DES}(\sigma)}$$

What subsets of permutations produce symmetric functions?

Which are Schur positive?

- Gessel: With fixed inverse descent set
- Gessel–Reutenaur: With fixed cycle type
- Elizalde–Roichman: Arc permutations
- Adin–Roichman: Closed under Knuth relations

Schur positive (classical) descent enumerators

For a subset $A \subseteq \mathfrak{S}_n$, define

$$\mathcal{Q}_A(\mathbf{x}) = \sum_{\pi \in A} F_{n, \text{DES}(\sigma)}$$

What subsets of permutations produce symmetric functions?

Which are Schur positive?

- Gessel: With fixed inverse descent set
- Gessel–Reutenaur: With fixed cycle type
- Elizalde–Roichman: Arc permutations
- Adin–Roichman: Closed under Knuth relations
- Hamaker–Pawłowski–Sagan: Avoiding certain collections of length 3 patterns

Schur positive (classical) descent enumerators

For a subset $A \subseteq \mathfrak{S}_n$, define

$$\mathcal{Q}_A(\mathbf{x}) = \sum_{\pi \in A} F_{n, \text{DES}(\sigma)}$$

What subsets of permutations produce symmetric functions?

Which are Schur positive?

- Gessel: With fixed inverse descent set
- Gessel–Reutenaur: With fixed cycle type
- Elizalde–Roichman: Arc permutations
- Adin–Roichman: Closed under Knuth relations
- Hamaker–Pawłowski–Sagan: Avoiding certain collections of length 3 patterns
- Many more

Schur positive small descent enumerators

For a subset $A \subseteq \mathfrak{S}_n$, define

$$SQ_A(\mathbf{x}) = \sum_{\sigma \in A} F_{n, \text{SDES}(\sigma)}$$

What subsets of permutations produce symmetric functions?
Which are Schur positive?

Schur positive small descent enumerators

For a subset $A \subseteq \mathfrak{S}_n$, define

$$SQ_A(\mathbf{x}) = \sum_{\sigma \in A} F_{n, \text{SDES}(\sigma)}$$

What subsets of permutations produce symmetric functions?

Which are Schur positive?

- Chow, Grinberg–Stanley: All permutations

Schur positive small descent enumerators

For a subset $A \subseteq \mathfrak{S}_n$, define

$$SQ_A(\mathbf{x}) = \sum_{\sigma \in A} F_{n, \text{SDES}(\sigma)}$$

What subsets of permutations produce symmetric functions?
Which are Schur positive?

- Chow, Grinberg–Stanley: All permutations
- That's it as far as previous results go!

Schur positive small descent enumerators

For a subset $A \subseteq \mathfrak{S}_n$, define

$$SQ_A(\mathbf{x}) = \sum_{\sigma \in A} F_{n, \text{SDES}(\sigma)}$$

What subsets of permutations produce symmetric functions?
Which are Schur positive?

- Chow, Grinberg–Stanley: All permutations
- That's it as far as previous results go!

Corollary (C. +26)

Let Π be a finite set of patterns such that $\text{SDES}(\pi) = \emptyset$ for all $\pi \in \Pi$. Then, $SQ_{A_{V_n}(\Pi)}(\mathbf{x})$ is symmetric and Schur positive.

Schur positive small descent enumerators

For a subset $A \subseteq \mathfrak{S}_n$, define

$$SQ_A(\mathbf{x}) = \sum_{\sigma \in A} F_{n, \text{SDES}(\sigma)}$$

What subsets of permutations produce symmetric functions?
Which are Schur positive?

- Chow, Grinberg–Stanley: All permutations
- That's it as far as previous results go!

Corollary (C. +26)

Let Π be a finite set of patterns such that $\text{SDES}(\pi) = \emptyset$ for all $\pi \in \Pi$. Then, $SQ_{A_{V_n}(\Pi)}(\mathbf{x})$ is symmetric and Schur positive.

This is implied by the theorem stated earlier.

Example

Let $A = \{2341, 2413, 3142, 3421, 4123, 4312\}$
(all permutations of cycle type (4))

$$Q_A = s_{2,1,1} + s_{3,1} \quad \mathcal{SQ}_A = F_{4,\{1\}} + F_{4,\{3\}} + 4F_{4,\{\emptyset\}}$$

Let $B = \{123, 213, 132, 312, 321\} = \text{Av}_3(231)$

$$Q_B = F_{3,\{1,2\}} + 2F_{3,\{2\}} + F_{3,\{1\}} + F_{3,\emptyset} \quad \mathcal{SQ}_B = s_{1,1,1} + s_{2,1} + 2s_3$$

Example

Let $A = \{2341, 2413, 3142, 3421, 4123, 4312\}$
(all permutations of cycle type (4))

$$Q_A = s_{2,1,1} + s_{3,1} \quad \mathcal{S}Q_A = F_{4,\{1\}} + F_{4,\{3\}} + 4F_{4,\{\emptyset\}}$$

Let $B = \{123, 213, 132, 312, 321\} = \text{Av}_3(231)$

$$Q_B = F_{3,\{1,2\}} + 2F_{3,\{2\}} + F_{3,\{1\}} + F_{3,\emptyset} \quad \mathcal{S}Q_B = s_{1,1,1} + s_{2,1} + 2s_3$$

Conclusion: Q_A and $\mathcal{S}Q_A$ are very different

Using the following symmetric function identity of Gessel

$$s_{(n-i,1^i)} = \sum_{\substack{S \subseteq [n-1] \\ |S|=i}} F_{n,S},$$

if $a_k^\Pi = |\{\sigma \in \text{Av}_k(\Pi) : \text{SDES}(\sigma) = \emptyset\}|$, then we see

$$\begin{aligned} \sum_{\sigma \in \text{Av}_n(\Pi)} F_{n, \text{SDES}(\sigma)} &= \sum_{S \subseteq [n-1]} F_{n,S} |\{\sigma \in \text{Av}_n(\Pi) : \text{SDES}(\sigma) = S\}| \\ &= \sum_{S \subseteq [n-1]} a_{n-|S|}^\Pi F_{n,S} \\ &= \sum_{i=0}^{n-1} a_{n-i}^\Pi \sum_{\substack{S \subseteq [n-1] \\ |S|=i}} F_{n,S} \\ &= \sum_{i=0}^{n-1} a_{n-i}^\Pi s_{(n-i,1^i)}. \end{aligned}$$



Refined expansion

Let $\sigma \in \mathfrak{S}_n$ be a permutation and set $\text{des}(\sigma) = |\text{DES}(\sigma)|$ and $\text{sdes}(\sigma) = |\text{SDES}(\sigma)|$. Also set $\text{asc}(\sigma) = n - 1 - \text{des}(\sigma)$ i.e. the number of **ascents** of σ .

Corollary (C. +26)

If Π is a set of patterns such that $\text{sdes}(\pi) = 0$ for all $\pi \in \Pi$, then for all $n \geq 1$, we have

$$\sum_{\sigma \in \text{Av}_n(\Pi)} q^{\text{des}(\sigma) - \text{sdes}(\sigma)} t^{\text{asc}(\sigma)} F_{n, \text{SDES}(\sigma)} = \sum_{i=1}^n a_i^{\Pi}(q, t) s_{i, 1^{n-i}}$$

where

$$a_i^{\Pi}(q, t) = \sum_{\substack{\sigma \in \text{Av}_i(\Pi) \\ \text{sdes}(\sigma)=0}} q^{\text{des}(\sigma)} t^{\text{asc}(\sigma)}.$$

Refined Expansion


The coefficients $a_i^\Pi(q, t)$ of $s_{i, 1^{n-i}}$ do not depend on n .

Example ($\Pi = \{231\}$)


- ① s_1
- ② $s_{1,1} + ts_2$
- ③ $s_{1,1,1} + ts_{2,1} + (qt + t^2) s_3$
- ④ $s_{1,1,1,1} + ts_{2,1,1} + (qt + t^2) s_{3,1} + (3qt^2 + t^3) s_4$
- ⑤ $s_{1,1,1,1,1} + ts_{2,1,1,1} + (qt + t^2) s_{3,1,1} + (3qt^2 + t^3) s_{4,1} + (2q^2t^2 + 6qt^3 + t^4) s_5$
- ⑥ $s_{1,1,1,1,1,1} + ts_{2,1,1,1,1} + (qt + t^2) s_{3,1,1,1} + (3qt^2 + t^3) s_{4,1,1} + (2q^2t^2 + 6qt^3 + t^4) s_{5,1} + (10q^2t^3 + 10qt^4 + t^5) s_6$
- ⑦ $s_{1,1,1,1,1,1,1} + ts_{2,1,1,1,1,1} + (qt + t^2) s_{3,1,1,1,1} + (3qt^2 + t^3) s_{4,1,1,1} + (2q^2t^2 + 6qt^3 + t^4) s_{5,1,1} + (10q^2t^3 + 10qt^4 + t^5) s_{6,1} + (5q^3t^3 + 30q^2t^4 + 15qt^5 + t^6) s_7$

Refined expansion–Proof


The earlier bijection preserves **big descents** and **ascents**



(a) σ



(b) u (spread out)



(c) u



Coefficient polynomials


What are the coefficient polynomials?

$$a_i^\Pi(q, t) = \sum_{\substack{\sigma \in \text{Av}_i(\Pi) \\ \text{sdes}(\sigma)=0}} q^{\text{des}(\sigma)} t^{\text{asc}(\sigma)}.$$

Coefficient polynomials

What are the coefficient polynomials?

$$a_i^\Pi(q, t) = \sum_{\substack{\sigma \in \text{Av}_i(\Pi) \\ \text{sdes}(\sigma)=0}} q^{\text{des}(\sigma)} t^{\text{asc}(\sigma)}.$$



Motzkin path $P \in M_8$

Theorem

Let $\#D(P)$ and $\#F(P)$ be set of down steps and flat steps of a Motzkin path P . Then for all $i \geq 1$,

$$a_i^{231}(q, t) = \sum_{\substack{\sigma \in \text{Av}_i(\Pi) \\ \text{sdes}(\sigma)=0}} q^{\text{des}(\sigma)} t^{\text{asc}(\sigma)} = \sum_{P \in M_{i-1}} q^{\#D(P)} t^{\#F(P)+\#D(P)}$$

Switching gears

For a finite subset $A \subseteq \mathbb{Z}_{>0}^n$, define

$$\mathcal{Q}_A(\mathbf{x}) = \sum_{\sigma \in A} F_{n, \text{DES}(\sigma)}$$

What set of **words** produce symmetric functions? Which are Schur positive?

Switching gears

For a finite subset $A \subseteq \mathbb{Z}_{>0}^n$, define

$$Q_A(\mathbf{x}) = \sum_{\sigma \in A} F_{n, \text{DES}(\sigma)}$$

What set of **words** produce symmetric functions? Which are Schur positive?

For example, if A any set of words that is invariant under the standard action of $\mathfrak{S}_n$ on $\mathbb{Z}_{>0}^n$, then $Q_A(\mathbf{x})$ is Schur positive.

Switching gears

For a finite subset $A \subseteq \mathbb{Z}_{>0}^n$, define

$$Q_A(\mathbf{x}) = \sum_{\sigma \in A} F_{n, \text{DES}(\sigma)}$$

What set of **words** produce symmetric functions? Which are Schur positive?

For example, if A any set of words that is invariant under the standard action of $\mathfrak{S}_n$ on $\mathbb{Z}_{>0}^n$, then $Q_A(\mathbf{x})$ is Schur positive.

Not true for $\mathcal{S}Q_A(\mathbf{x})$: $\mathcal{S}Q_{\mathfrak{S}_4 \cdot \{1123\}} = s_{2,1,1} - s_{2,2} + 2s_{3,1} + 5s_4$.

Switching gears

For a finite subset $A \subseteq \mathbb{Z}_{>0}^n$, define

$$Q_A(\mathbf{x}) = \sum_{\sigma \in A} F_{n, \text{DES}(\sigma)}$$

What set of **words** produce symmetric functions? Which are Schur positive?

For example, if A any set of words that is invariant under the standard action of $\mathfrak{S}_n$ on $\mathbb{Z}_{>0}^n$, then $Q_A(\mathbf{x})$ is Schur positive.

Not true for $\mathcal{S}Q_A(\mathbf{x})$: $\mathcal{S}Q_{\mathfrak{S}_4 \cdot \{1123\}} = s_{2,1,1} - s_{2,2} + 2s_{3,1} + 5s_4$.

We will see that $\mathcal{S}Q_A(\mathbf{x})$ is Schur positive for the set of *parking functions*.

Parking functions

A **parking function** of length n is a word $w = w_1 \cdots w_n$ (i.e. a function $w : [n] \rightarrow \mathbb{Z}_{>0}$) such that $|w^{-1}([i])| \geq i$ for all i . Let PF_n denote the set of parking functions of length n .

$$\text{PF}_3 = \{111, 112, 121, 211, 113, 131, 311, 122, 212, 221, \\ 123, 132, 213, 231, 312, 321\}$$

Parking Functions

- Let $\text{sasc}(w)$ be the number of $i \in [n-1]$ such that $w_i = w_{i+1} - 1$ (**small ascents**)
- Let $\text{tie}(w)$ be the number of $i \in [n-1]$ such that $w_i = w_{i+1}$ (**ties**)

Theorem

For all $n \geq 1$,

$$\sum_{w \in \text{PF}_n} q^{\text{sasc}(w)} t^{\text{tie}(w)} F_{n, \text{SDES}(w)} = \sum_{i=1}^n (q + t + n - 2)^{i-1} s_{i, 1^{n-i}}$$

Proof involves passing to “Prüfer codes.”

Parking Functions

The Prüfer code proof actually reveals a surprising symmetry

Theorem

For all $n \geq 1$, any permutation of the statistics defines the same, Schur positive function

$$\sum_{i=1}^n (q+t+n-2)^{i-1} s_{i,1^{n-i}} = \begin{cases} \sum_{w \in \text{PF}_n} q^{\text{sasc}(w)} t^{\text{tie}(w)} F_{n, \text{SDES}(w)} \\ \sum_{w \in \text{PF}_n} q^{\text{tie}(w)} t^{\text{sasc}(w)} F_{n, \text{SDES}(w)} \\ \sum_{w \in \text{PF}_n} q^{\text{sdes}(w)} t^{\text{sasc}(w)} F_{n, \text{TIE}(w)} \\ \vdots \end{cases}$$

Let $SQ_{\text{PF}_n}(q, t)$ denote this common value.

Expansion in the p (for parking function) basis

For a partition $\lambda = (1^{m_1} 2^{m_2} \dots)$, let $z_\lambda = 1^{m_1} m_1! 2^{m_2} m_2! \dots$

Theorem

Let y be an indeterminate. Then for all $n \geq 1$,

$$\sum_{i=1}^n y^{i-1} s_{i, 1^{n-i}} = \frac{1}{y+1} \sum_{\lambda \vdash n} \frac{p_\lambda}{z_\lambda} \prod_{i=1}^{\ell(\lambda)} (y^{\lambda_i} - (-1)^{\lambda_i}).$$

Applies the Adin–Roichman s -to- p formula to hook shapes

Corollary

For all $n \geq 1$,

$$\mathcal{SQ}_{\text{PF}_n}(q, t) = \frac{1}{q+t+n-1} \sum_{\lambda \vdash n} \frac{p_\lambda}{z_\lambda} \prod_{i=1}^{\ell(\lambda)} ((q+t+n-2)^{\lambda_i} - (-1)^{\lambda_i})$$

- Our result

$$\sum_{\sigma \in \text{Av}_n(\Pi)} q^{\text{des}(\sigma) - \text{sdes}(\sigma)} t^{\text{asc}(\sigma)} F_{n, \text{SDES}(\sigma)} = \sum_{i=1}^n a_i^{\Pi}(q, t) s_{i, 1^{n-i}}$$

says that there is an action of $\mathfrak{S}_n$ on $\text{Av}_n(\Pi)$ that preserves $(\text{des}(\sigma) - \text{sdes}(\sigma), \text{asc}(\sigma))$. It would be very interesting to actually construct it.

- Our result

$$\sum_{\sigma \in \text{Av}_n(\Pi)} q^{\text{des}(\sigma) - \text{sdes}(\sigma)} t^{\text{asc}(\sigma)} F_{n, \text{SDES}(\sigma)} = \sum_{i=1}^n a_i^\Pi(q, t) s_{i, 1^{n-i}}$$

says that there is an action of $\mathfrak{S}_n$ on $\text{Av}_n(\Pi)$ that preserves $(\text{des}(\sigma) - \text{sdes}(\sigma), \text{asc}(\sigma))$. It would be very interesting to actually construct it.

- $a_i^{123}(1, 1) = a_i^{231}(1, 1)$, but $a_i^{123}(q, t) \neq a_i^{231}(q, t)$. What statistics on Motzkin paths does $a_i^{123}(q, t)$ enumerate?

- Our result

$$\sum_{\sigma \in \text{Av}_n(\Pi)} q^{\text{des}(\sigma) - \text{sdes}(\sigma)} t^{\text{asc}(\sigma)} F_{n, \text{SDES}(\sigma)} = \sum_{i=1}^n a_i^\Pi(q, t) s_{i, 1^{n-i}}$$

says that there is an action of $\mathfrak{S}_n$ on $\text{Av}_n(\Pi)$ that preserves $(\text{des}(\sigma) - \text{sdes}(\sigma), \text{asc}(\sigma))$. It would be very interesting to actually construct it.

- $a_i^{123}(1, 1) = a_i^{231}(1, 1)$, but $a_i^{123}(q, t) \neq a_i^{231}(q, t)$. What statistics on Motzkin paths does $a_i^{123}(q, t)$ enumerate?
- More generally, look for how small descents fit into the broader place that descents have in combinatorics (e.g. Solomon's descent algebra)

- Our result

$$\sum_{\sigma \in \text{Av}_n(\Pi)} q^{\text{des}(\sigma) - \text{sdes}(\sigma)} t^{\text{asc}(\sigma)} F_{n, \text{SDES}(\sigma)} = \sum_{i=1}^n a_i^\Pi(q, t) s_{i, 1^{n-i}}$$

says that there is an action of $\mathfrak{S}_n$ on $\text{Av}_n(\Pi)$ that preserves $(\text{des}(\sigma) - \text{sdes}(\sigma), \text{asc}(\sigma))$. It would be very interesting to actually construct it.

- $a_i^{123}(1, 1) = a_i^{231}(1, 1)$, but $a_i^{123}(q, t) \neq a_i^{231}(q, t)$. What statistics on Motzkin paths does $a_i^{123}(q, t)$ enumerate?
- More generally, look for how small descents fit into the broader place that descents have in combinatorics (e.g. Solomon's descent algebra)

Thanks!